



## Coupling the endophytic microbiome with the host transcriptome in olive roots



Antonio J. Fernández-González<sup>a</sup>, Jorge A. Ramírez-Tejero<sup>b</sup>, María Patricia Nevado-Berzosa<sup>b</sup>, Francisco Luque<sup>b</sup>, Manuel Fernández-López<sup>a</sup>, Jesús Mercado-Blanco<sup>c,\*</sup>

<sup>a</sup> Departamento de Microbiología del Suelo y Sistemas Simbióticos, Estación Experimental del Zaidín, Consejo Superior de Investigaciones Científicas (CSIC), Calle Profesor Albareda 1, 18008 Granada, Spain

<sup>b</sup> Departamento de Biología Experimental, Centro de Estudios Avanzados en Olivar y Aceites de Oliva, Universidad de Jaén, Jaén 23071, Spain

<sup>c</sup> Departamento de Protección de Cultivos, Instituto de Agricultura Sostenible, CSIC, Campus 'Alameda del Obispo' s/n, Avd. Menéndez Pidal s/n, 14004 Córdoba, Spain

### ARTICLE INFO

#### Article history:

Received 13 May 2021

Received in revised form 6 August 2021

Accepted 20 August 2021

Available online 21 August 2021

#### Keywords:

*Actinophytocola* spp.

Holobiont

*Olea europaea* L.

Olive transcriptome

Root endophytome

Stress tolerance

### ABSTRACT

The connection between olive genetic responses to environmental and agro-climatic conditions and the composition, structure and functioning of host-associated, belowground microbiota has never been studied under the holobiont conceptual framework. Two groups of cultivars growing under the same environmental, pedological and agronomic conditions, and showing highest (AH) and lowest (AL) *Actinophytocola* relative abundances, were earlier identified. We aimed now to: i) compare the root transcriptome profiles of these two groups harboring significantly different relative abundances in the above-mentioned bacterial genus; ii) examine their rhizosphere and root-endosphere microbiota co-occurrence networks; and iii) connect the root host transcriptome pattern to the composition of the root microbial communities by correlation and co-occurrence network analyses. Significant differences in olive gene expression were found between the two groups. Co-occurrence networks of the root endosphere microbiota were clearly different as well. Pearson's correlation analysis enabled a first portrayal of the interaction occurring between the root host transcriptome and the endophytic community. To further identify keystone operational taxonomic units (OTUs) and genes, subsequent co-occurrence network analysis showed significant interactions between 32 differentially expressed genes (DEGs) and 19 OTUs. Overall, negative correlation was detected between all upregulated genes in the AH group and all OTUs except of *Actinophytocola*. While two groups of olive cultivars grown under the same conditions showed significantly different microbial profiles, the most remarkable finding was to unveil a strong correlation between these profiles and the differential gene expression pattern of each group. In conclusion, this study shows a holistic view of the plant-microbiome communication.

© 2021 The Authors. Published by Elsevier B.V. on behalf of Research Network of Computational and Structural Biotechnology. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

The olive tree (*Olea europaea* L.) is iconic in the Mediterranean Basin and has strong economic, social and ecological impacts in this region [1]. While the origin and domestication of this crop has not yet been undoubtedly documented, it is likely that took place in the Levant area from where olive cultivation gradually expanded along with human migrations [2,3]. Multilocal selection and backcrosses between wild (*O. europaea* L. subsp. *europaea* var.

*sylvestris* (Mill.) Leh.) and cultivated (*O. europaea* L. subsp. *europaea* var. *europaea*) olives have created the huge number of cultivars currently available [4]. Around 10.5 million ha are now dedicated to olive cultivation worldwide, creating an agro-ecosystem of utmost relevance in countries of the Mediterranean Basin where 98% of the cultivated area is located [5]. The major product of this tree crop is virgin olive oil, a source of fat providing nutritional and human health benefits that include reduction in the prevalence of cardiovascular diseases, prevention of some types of cancers and type 2 diabetes, and stimulation of higher diversity of beneficial gut bacteria [6–8].

In contrast to most of the fruit crops, the olive tree is well adapted to environmental conditions usually found in Mediterranean-type climatic regions (e.g. wide thermal amplitude,

\* Corresponding author.

E-mail addresses: [antonio.fernandez@eez.csic.es](mailto:antonio.fernandez@eez.csic.es) (A.J. Fernández-González), [jrtejero@ujaen.es](mailto:jrtejero@ujaen.es) (J.A. Ramírez-Tejero), [mnevado@ujaen.es](mailto:mnevado@ujaen.es) (M.P. Nevado-Berzosa), [flluque@ujaen.es](mailto:flluque@ujaen.es) (F. Luque), [manuel.fernandez@eez.csic.es](mailto:manuel.fernandez@eez.csic.es) (M. Fernández-López), [jesus.mercado@ias.csic.es](mailto:jesus.mercado@ias.csic.es) (J. Mercado-Blanco).

salt stress, water deprivation, etc.) [9,10]. Tolerance to some of these conditions (i.e. drought and/or saline soils) appears to be genotype dependent although studies are only limited to few cultivars [11–13]. While the plant genotype is clearly decisive to explain adaptation to diverse environmental situations, the contribution of its associated microbiota is just starting to be unveiled. This is particularly true for the plant root/belowground microbiota pair since crucial multitrophic interactions for the development, fitness and health of the plant holobiont take place in the rhizosphere/root endosphere compartments [14]. However, the connection between the global host genetic responses to different environmental and agro-climatic conditions and the composition, structure and functioning of its inhabiting microbiota remains mostly unknown. While pedological characteristics, plant age and phenology, variable weather conditions and root exudates [15–17] are relevant, the plant genotype often appears as the most decisive factor to shape the composition and functioning of the belowground microbiome, particularly at the root endosphere level [18–23].

Our knowledge about the link between olive-associated microbial communities and growth, development, fitness and adaptation to a/biotic constraints of this long-living tree is still very fragmentary. Nevertheless, recent studies are providing significant information about the factors shaping the olive-associated microbiota and how their constituents contribute to host stress tolerance and adaptation to specific soil/climatic conditions. For instance, different soil management practices affected the composition of the arbuscular mycorrhizal fungi (AMF) community [24]. The olive endophytic bacteriome has been shown to originate from the soil, reaching aerial organs through the xylem sap, and the composition of these communities seemed to be largely influenced by agronomic practices implemented in the orchard [25,26]. Our own studies have demonstrated that agricultural management produces differences in the belowground microbiota compared to that in adjacent soils left under natural conditions. Agronomic practices also removed differences in the microbial communities associated with olive cultivars [27]. Other works have shown that olive cultivars played a more decisive structuring effect than plant organs on the composition of the epiphytic and endophytic bacterial communities of the phyllosphere [28]. Most of the available studies on the olive microbiota are related to biotic constraints caused by pathogens and arthropods such as *Xylella fastidiosa*, causing the olive quick decline syndrome [29–31], *Pseudomonas savastanoi* pv. *savastanoi*, the etiological agent of the olive knot disease [32], or *Bactrocera oleae*, the olive fruit fly [33]. Our own studies have revealed that the presence of the soil-borne fungus *Verticillium dahliae*, causing one of the most serious disease affecting olive cultivation (Verticillium wilt) [34], did not produce relevant alterations in the structure and functionality of the belowground microbiota of two olive cultivars differing in susceptibility to the disease. Remarkably, however, notable differences were found in their co-occurrence networks in response to the inoculation with the pathogen, particularly in the root endosphere communities [35].

Concerning the host side of the olive holobiont, the most recent whole transcriptome-based studies are providing interesting information on aspects such as how and where the domestication process might have taken place [36], the basis of the alternate olive fruit bearing [37], the complex gene expression patterns in different organs [38], the genetics controlling drupe maturation and response to environmental cues [39], or the development process from germination to the juvenile stage [40]. Likewise, global genetic responses to different stressing situations have been studied using transcriptomic approaches as for the case of olive adaptation to cold temperatures [41,42] or salinity [43]. RNA-seq studies have been very useful to unravel the differential responses of olive cultivars upon the attack of pathogens such as *X. fastidiosa*

[44] or *V. dahliae* [45]. Moreover, dual or co-transcriptomics approaches offered the opportunity to unveil the interaction between olive and *V. dahliae* at full transcriptome level and during the interaction between both partners [46,47]. However, the cross-talk between olive and its inhabiting microbiota has not been studied from a holistic perspective. Recently, available RNA-seq datasets from root tissues of *V. dahliae*-infected and non-infected plants have been re-examined in an attempt to describe the disease through a systems biology approach [48]. While data mining from RNA-seq studies offers a novel and interesting perspective, the role of important components of the plant-associated microbiota to explain the reaction to a specific stress situation can be overlooked to some degree due to the methodological approaches implemented [48,49].

We recently described the belowground microbiota (rhizosphere and root endosphere) associated to different olive cultivars grown under the same climatic, agronomical and pedological conditions [20]. Overall, results showed that the olive genotype was the main factor shaping the root-associated microbial communities, and that the olive root endosphere microbiota was less diverse than that of the rhizosphere. Remarkably, *Actinophytocola* (*Actinobacteria*, *Pseudonocardiaceae* family) was found to be the most abundant bacterial genus in the olive root endosphere, suggesting it may play an important role in olive adaptation to environmental stresses. Indeed, representatives of *Actinophytocola* spp. have been found in highly arid environments like the Sahara, Tenger Desert (China) and Mongolia [50–53]. Interestingly, *Actinophytocola* was not only isolated from soils but also from inside the root of Thai glutinous rice plants (*Oryza sativa* sp.) [54].

Considering the previous antecedents, and to further explore the meaning of the significant differences in *Actinophytocola* relative abundance previously found in roots of different olive cultivars, we aimed in this study to: i) compare the olive root gene expression, at the whole transcriptome level, of two groups of olive cultivars harboring significantly-different abundances of the genus *Actinophytocola*; ii) analyze the rhizosphere and root-endosphere co-occurrence networks of the microbial communities associated to these two groups; and iii) connect the host whole-transcriptome expression to the composition of the root residing microbial communities by correlation and co-occurrence network (combining host relative gene expression and operational taxonomic units (OTUs) relative abundance data) analyses. The hypothesis to-be-tested is that diverse olive varieties cultivated under the same (agronomic, climatic, pedological) conditions show different root transcriptome patterns, and that these global gene expression differences can affect keystone OTUs (e.g. *Actinophytocola*) of the root endophytome that might be relevant for the fitness of the olive holobiont. This global interaction can be demonstrated by implementing an approach involving host transcriptomics and microbiomics.

## 2. Methods

### 2.1. Starting point data

The study of Fernández-González and co-workers [20], in which the microbial communities of the root system of 36 cultivars from the World Olive Germplasm Collection (WOGC) were studied, was taken as a starting point, using the same OTUs tables. In addition, transcriptomics data published by Ramírez-Tejero and co-workers [55] were also considered. Thus, the trees surveyed in these two previous studies constituted the biological samples (individuals) considered for this present study. In fact, both soil and root samples for metabarcoding and transcriptomic analysis originated from the same sampling event. Indeed, all samples were

collected on April 26th, 2017 during the flowering period, after a month without rainfall (Additional File 1: Table S1) or irrigation. The only difference between the two studies was that for the metabarcoding study three replicates were taken per cultivar, while for transcriptomics there were two replicates. Therefore, the two replicates that matched in both studies were the ones used in the correlation analyses between genes and OTUs.

## 2.2. Selection of olive cultivars according to the differential relative abundance of *Actinophytocola*

Of the 36 cultivars previously mentioned, 10 were selected for this study because of their differences on relative abundance of the bacterial genus *Actinophytocola* previously reported [20]. We particularly focused on differences found in the root endosphere compartment. Thus, 'Koroneiki', 'Mavreya', 'Myrtolia', 'Uslu' and 'Verdial de Vélez-Málaga' were preliminary selected as the extreme representatives of the *Actinophytocola* high-prevalence (AH) group while 'Barri', 'Klon\_14\_1812', 'Maarri', 'Mastoidis' and 'Shengeh' were chosen as the extreme members of the *Actinophytocola* low-prevalence (AL) group.

## 2.3. Gene expression analysis

To compare gene expression profiles between the two groups of cultivars, AH and AL, the gene expression study was conducted using the DNASTar (ArrayStar 16) Qseq software for RNA-seq analyses (www.dnastar.com). PE Illumina reads were mapped using as reference the complete transcriptome of *Olea europaea* cv. Picual [56]. Mapping was undertaken using high-stringency parameters k-mer = 63 and 95% of matches. The default normalization method of reads per kilobase per million mapped (RPKM) was implemented. Results and associated charts were processed with Microsoft Excel. A fold change (FC)  $\geq 8$  and a *p*-value  $\leq 0.01$  were considered to detect differentially expressed genes (DEGs) between AH and AL. Thus, comparisons between groups consisted in 10 biological replicates of each group, corresponding to five cultivars and two trees from each one.

## 2.4. Statistical analyses

To assess the differences in *Actinophytocola* relative abundance between the AH and AL groups, both groups were compared by the Student's *t*-test. In addition,  $\alpha$ -diversity indices (observed richness; Shannon and inverse of Simpson diversity; and Shannon evenness) were compared with the Mann-Whitney-Wilcoxon's *U* test using the R function 'wilcox.test'. These comparisons were split and the number of sequences was rarefied by compartment (i.e. root endosphere and rhizosphere) and by domain (i.e. bacteria and fungi). To analyze  $\beta$ -diversity, a normalization of the filtered OTUs sequence counts was performed using the 'trimmed means of M' (TMM) method with the BioConductor package 'edgeR' [57]. The normalized data were considered to perform Non-Metric MultiDimensional Scaling (NMDS) on Bray-Curtis dissimilarities to ordinate in two dimensions the variance of  $\beta$ -diversity between the AH and AL groups. Similarly as for  $\alpha$ -diversity, these comparisons were split by compartment and by domain. Ordination analyses were performed using the R package 'phyloseq' [58]. To analyze the effect of the AH versus AL grouping on community dissimilarities, permutational analysis of variance (PERMANOVA) and permutational analysis of multivariate dispersions (BETADISPER) were performed using the functions 'adonis' and 'betadisper' in the 'vegan package' with 9,999 permutations [59]. All of the above-mentioned analyses that were carried out using R have been provided in a script (Additional File 2: Statistical\_analyses\_script.R).

## 2.5. Pearson's correlations and heatmap construction

In order to perform the correlation analysis between DEGs and microbial OTUs in the root endosphere of all cultivars under study (i.e. AH and AL groups together), gene expression values (reads per kilobase per million mapped reads; RPKM) were multiplied by 1000. Therefore, setting the smallest value, bigger than 0, to 1 in the dataset. Then, the replicate from which only microbial but not gene information was available was removed (see section 2.1). Eventually, two replicates were kept for each cultivar. Thus, cultivar 'Shengeh' was excluded due to the low number of sequences in the fungal community of this cultivar ( $n = 18$ ). Subsequently, the data were log-transformed to be able to compare them. To eliminate spurious correlations, only those OTUs found in at least half of the samples were retained ( $n = 308$  OTUs in  $n = 9$  replicates). The correlation between DEGs and OTUs was made with the Pearson correlation coefficient using the 'corrplot' [60] package and a heatmap was constructed with the 'pheatmap' [61] package, both available in R. Pearson's correlation coefficient cut-off of 0.6 and *p*-value  $< 0.05$  were established to consider significant and strong correlations. To sort the heatmap, hierarchical clustering dendrograms were carried out for DEGs and OTUs with a complete linkage clustering method and using Euclidean distances with the software MultiExperimentViewer v.4.9.0 [62].

## 2.6. Network construction, comparison and visualization

Microbial (bacterial and fungal) co-occurrence networks were separately constructed for each group (AH and AL) and each compartment (root endosphere and rhizosphere) by using MENAP [63] and according to our previous implementation [35]. Three replicates were used per cultivar ( $n = 15$ , see section 2.1), except of AL endosphere, for which "Shengeh" had to be removed because insufficient fungal reads. Moreover, 100 random networks were performed to each empirical network for comparing the major topological properties between groups with Student's *T* tests, as earlier described [64–66]. All the networks were visualized by using Cytoscape v.3.7.1 [67]. The node properties tables were analyzed in Microsoft Excel to plot and highlight the keystone OTUs according to their within-module connections ( $Z_i$ ) and among-module connections ( $P_i$ ) values.

To generate the co-occurrence network between DEGs and OTUs, the log-transformed and filtered dataset previously obtained to construct the heatmap were used (see section 2.5). By using Pearson's correlation coefficient, a network with an *R* square of power-law value of 0.861 was obtained; that is, it gave rise to a robust and far from random network. A lower value was obtained when building the network using the Spearman's correlation coefficient. Due to this result Pearson was used as the best correlation method in both the heatmap and the co-occurrence network. The network construction method of MENAP web software is based on the comparison of the correlation matrix against an RMT (Random Matrix Theory) model. In summary, the RMT-based method informs whether correlations of the matrix (between OTUs and DEGs in our study) follow a Gaussian distribution, confirming a true correlation and discarding a Poisson distribution suggestive of an artefactual correlation. Eventually, this methodological approach predicts which cut-off value for Pearson is the ideal one to filter out all non-significant correlations (parameter that in other methods of network construction must be chosen by the user). In our case, the algorithm determined that the most appropriate cut-off Pearson's value was 0.8, and not 0.6 which was used as a cut-off value to consider a strong correlation when constructing the heatmap (section 2.5).

### 3. Results

#### 3.1. Differential distribution of endophytic *Actinophytocola* among olive cultivars

We earlier described the composition of the bacterial and fungal endophytic communities inhabiting olive roots from 36 cultivars comprising a subset of a so-called “olive core collection” [20]. Remarkably, data showed that the bacterial genus *Actinophytocola* (and others like *Flavitalea* although to a lesser extent) was found, by far, at significantly higher relative abundance in the olive root endosphere, with differences among cultivars though [20]. From this starting point, and in order to assess whether the observed significant differences in *Actinophytocola* relative abundance among olive cultivars correlate with differential host transcriptome responses, two groups (five cultivars each) displaying the highest and the lowest *Actinophytocola* relative abundances were made *ad hoc*. The relative abundance of *Actinophytocola* was determined by the proportional number of sequences for each cultivar replicate in the total number of bacterial reads. Five cultivars with the highest *Actinophytocola* prevalence constituted the AH group while five cultivars with the lowest *Actinophytocola* prevalence formed the AL group. The AH group had an average relative abundance of *Actinophytocola* of  $42.38 \pm 4.52$  (mean percentage  $\pm$  SE). In contrast, the AL group showed a mean prevalence of  $6.01 \pm 0.86$  (*t*-test *p*-value =  $1.30 \times 10^{-08}$ ). The AH group included cultivars Uslu (Turkey), Verdial de Vélez-Málaga (Spain), Koroneiki (Greece), Mavreya (Greece) and Myrtolia (Greece), while the AL group was comprised by cultivars Shengeh, (Iran), Mastoidis (Greece), Barri (Syria), Klon\_14\_1812 (Albania) and Maarri (Syria) (Fig. 1). No correlation was observed between the microbial communities of AH and AL cultivars and their geographical origins, nor with tolerance/susceptibility to biotic stresses such as *Verticillium* wilt [68] or Anthracnose (caused by *Colletotrichum* species) [69] (Fig. 1). Moreover, AH and AL cultivars distributed randomly between the two different phylogenies of cultivated olives recently reported [56].

#### 3.2. The AH and AL olive cultivars show significantly different whole-transcriptome profiles

In order to assess whether the *Actinophytocola* prevalence in the AH and AL groups was associated to different transcriptome

profiles in the host roots, the existence of DEGs among these groups was investigated. This analysis was performed using high stringency parameters to detect only highly consistent DEGs. As a result, differences in gene expression between the AH and AL groups at 8 FC and *p*-value  $\leq 0.01$  were found in 60 genes (Additional File 3: Fig. S1), 39 up-regulated in the AH group (Table 1 and Additional File 4: Fig. S2) and 21 up-regulated in the AL group (Table 2 and Additional File 5: Fig. S3). In the set of 39 up-regulated genes in the AH group and down-regulated in the AL cultivars three genes involved in plant defense to pathogens were found, one coding for a Germin-like protein subfamily 1 member 16 (Oleur061Scf2041g01041.1), and two *tlp* genes encoding Thaumatin-like proteins (Oleur061Scf0191g00025.1, Oleur061Scf0201g06012.1) with antifungal activity that are secreted to the apoplast. Additionally, one *CIGR1* gene (Oleur061Scf2742g00025.1) coding for a chitin-inducible gibberellin (GA)-responsive protein 1, that may play a regulatory role in the early step of oligosaccharide elicitor response [70], and a response-to-stress *ABA2* gene (Oleur061Scf2621g00015.1) encoding a zeaxanthin epoxidase, were also up-regulated in AH and down-regulated in AL cultivars. In addition, five genes (*ATL22*, *ATL46*, *ATL8*, *PP2B11* and *PCYOX1*; corresponding to Oleur061Scf0789g04009.1, Oleur061Scf4176g04002.1, Oleur061Scf1016g05039.1, Oleur061Scf7684g00004.1, Oleur061Scf3403g03008.1, respectively) involved in protein ubiquitination and/or protein degradation were also identified in this set of genes. In addition, four genes encoding proteins involved in protein biosynthesis pathway (Utp12, L34Ae, pre-mRNA-splicing factor 38 and protein translation factor SU11; corresponding to Oleur061Scf7124g01014.1, Oleur061Scf9068g00010.1, Oleur061Scf0172g00010.1, Oleur061Scf0579g00003.1, respectively) were also identified in this set, suggesting a higher protein turnover in AH roots. In this sense, gene expression was particularly activated in AH roots by two up-regulated transcription factors (TF), *HAT5* and *HOX6* (Oleur061Scf7420g01005.1, Oleur061Scf2236g05008.1), both coding for Homeobox-leucine zipper proteins. The *HOX6* gene may be involved in drought response [71]. In addition, a *XRN3* gene (Oleur061Scf1132g02001.1) coding for a suppressor of post-transcriptional gene silencing was up-regulated in AH roots as well. Another difference between AH and AL root transcriptomes was the expression of genes involved in the cell wall. Thus, in AH roots, two genes coding for Extensin proteins (Oleur061Scf3325g00004.1, Oleur061




Fig. 1. Relative abundance of the genus *Actinophytocola* in olive cultivars of the AL and AH groups. AH: *Actinophytocola* high-prevalence group; AL: *Actinophytocola* low-prevalence group. Tolerance (+) or susceptibility (-) of olive cultivars to *Verticillium dahliae* and *Colletotrichum* spp. are indicated. When this information was unavailable, cultivars are marked with “?”.



**Table 1**  
Upregulated genes in the *Actinophytocola* high-prevalence (AH) group of olive cultivars.

Gene code ID	Functional annotation	Relevant process or cellular compartment
Oleur061Scf0191g00025.1	Thaumatococcus-like protein IPR001938	Response to biotic stress. Antifungal activity. Secreted
Oleur061Scf0201g06012.1	Thaumatococcus-like protein IPR001938	Response to biotic stress. Antifungal activity. Secreted
Oleur061Scf2742g00025.1	Chitin-inducible gibberellin-responsive protein 1 IPR005202	Response to biotic stress
Oleur061Scf2041g01041.1	Germin-like protein subfamily 1 member 16 IPR001929	Response to biotic stress
Oleur061Scf2041g01034.1	Pentatricopeptide repeat-containing protein At3g29290 IPR002885	Response to biotic and abiotic stress
Oleur061Scf0789g04009.1	F-box protein PP2-B11 IPR025886	Protein turnover
Oleur061Scf4176g04002.1	RING-H2 finger protein ATL22 IPR013083	Protein turnover
Oleur061Scf1016g05039.1	RING-H2 finger protein ATL46 IPR013083	Protein turnover
Oleur061Scf7684g00004.1	RING-H2 finger protein ATL8 IPR013083	Protein turnover
Oleur061Scf3403g03008.1	Prenylcysteine oxidase IPR017046	Protein turnover
Oleur061Scf7124g01014.1	(Small-subunit processome, Utp12)	Protein synthesis
Oleur061Scf9068g00010.1	Protein translation factor SUI1 homolog IPR005874	Protein synthesis
Oleur061Scf0172g00010.1	(Ribosomal protein L34Ae)	Protein synthesis
Oleur061Scf0579g00003.1	Pre-mRNA-splicing factor 38A IPR005037	mRNA processing
Oleur061Scf1132g02001.1	5'-3' exoribonuclease 3 (XRN3) IPR001878	rRNA processing, miRNA degradation
Oleur061Scf3325g00004.1	(Extensin domain)	Cell wall component
Oleur061Scf2628g03012.1	(Pistil-specific extensin-like protein)	Cell wall component
Oleur061Scf3771g00037.1	Wall-associated receptor kinase 2 IPR000742	Cell wall component
Oleur061Scf1016g05003.1	(Six-bladed beta-propeller, TolB-like)	Cell wall (not clear)
Oleur061Scf7931g00024.1	Citrate synthase 3, peroxisomal IPR002020	Mitochondrial respiration
Oleur061Scf6638g01002.1	Protein FMP32, mitochondrial IPR024461	Mitochondrial respiration
Oleur061Scf3287g00010.1	Mitochondrial outer membrane porin IPR027246	Mitochondrial
Oleur061Scf6180g00005.1	(Probable transposase, PttA/En/Spm, plant)	DNA transposons
Oleur061Scf1200g04004.1	(Probable transposase, PttA/En/Spm, plant)	DNA transposons
Oleur061Scf1553g03011.1	(Sodium/calcium exchanger membrane region), IPR011992	Sodium/calcium exchanger
Oleur061Scf2621g00015.1	Zeaxanthin epoxidase, chloroplastic IPR008984 (SMAD/FHA domain),	Plant hormone biosynthesis (ABA)
Oleur061Scf8089g00019.1	L-ascorbate oxidase homolog IPR008972 (Cupredoxin)	Copper ion binding, oxidoreductase activity, secreted
Oleur061Scf1032g02009.1	B2 protein IPR013989	Development and cell death
Oleur061Scf3883g01014.1	Beta-glucuronosyltransferase GlcAT14B IPR003406	
Oleur061Scf7420g01005.1	Homeobox-leucine zipper protein HAT5 IPR003106	Transcription factor
Oleur061Scf2236g05008.1	Homeobox-leucine zipper protein HOX6 IPR003106	Transcription factor
Oleur061Scf2846g08008.1	(Protein of unknown function DUF597)	Transcription factor

Genes with higher expression in the AH group of cultivars at 8 FC and p-value  $\leq 0.01$ . Genes that may be involved in the same process or same compartment are represented with the same color. If processes are not the same but related, the same color with different tonality was used. Colorless means that the process is not known.

**Table 2**  
Upregulated genes in the *Actinophytocola* low-prevalence (AL) group of olive cultivars.

Gene code ID	Functional annotation	Relevant process or cellular compartment
Oleur061Scf1329g05010.1	Cytochrome P450 71A9 IPR001128	Response to biotic stress. Plant development
Oleur061Scf0889g00002.1	Cysteine-rich receptor-like protein kinase 3 IPR011009	Response to biotic stress. Plant development
Oleur061Scf9036g01047.1	Ethylene-responsive transcription factor ERF034 IPR016177	Response to biotic stress
Oleur061Scf1024g00024.1	Ethylene-responsive transcription factor 4 IPR016177	Response to biotic stress
Oleur061Scf0247g05018.1	E3 ubiquitin-protein ligase At3g02290 IPR013083	Protein turnover
Oleur061Scf2527g00016.1	Xyloglucan glycosyltransferase 4 IPR029044	Cell wall synthesis
Oleur061Scf5473g04020.1	DELLA protein GAI1 IPR005202	Plant growth and development. Root growth
Oleur061Scf3289g01014.1	Protein SHORT-ROOT IPR005202	Plant growth and development. Root growth
Oleur061Scf3112g01022.1	LRR receptor-like serine/threonine-protein kinase GSO1 IPR001611	Root growth
Oleur061Scf0888g00017.1	Polyol transporter 5 IPR005828	Sugar transporter
Oleur061Scf0276g01013.1	Glycerol-3-phosphate acyltransferase 1 IPR002123	Pollen development
Oleur061Scf0088g04051.1	LOB domain-containing protein 30 IPR004883	Xylem water transport
Oleur061Scf1251g02004.1	1-aminocyclopropane-1-carboxylate oxidase homolog IPR026992	Plant hormone biosynthesis (Ethylene)
Oleur061Scf0885g02003.1	Anthocyanidin 3-O-glucoside 6"-O-acyltransferase IPR003480	Pigment synthesis
Oleur061Scf1794g14034.1	Butyrate--CoA ligase AAE11, peroxisomal IPR000873	Lipid metabolism
Oleur061Scf3195g01012.1	Heparan-alpha-glucosaminide N-acetyltransferase	Lysosomal transport. Protein complex oligomerization
Oleur061Scf0724g05015.1	(DC1), IPR013083	
Oleur061Scf3229g02021.1	(Leucine-rich repeat domain, L domain-like)	
Oleur061Scf7440g00010.1	GTP-binding nuclear protein Ran/TC4 IPR001806	
Oleur061Scf0269g02013.1	Unknown protein	
Oleur061Scf0604g11007.1	Unknown protein	

Genes with higher expression in the AH group of cultivars at 8 FC and p-value  $\leq 0.01$ . Genes that may be involved in the same process or same compartment are represented with the same color. If processes are not the same but related, the same color with different tonality was used. Colorless means that the process is not known.

Scf2628g03012.1) that form a structural component of the primary cell wall and strengthen it, were found up-regulated. Related to this, a *WAK2* gene (Oleur061Scf3771g00037.1) encoding a wall-associated receptor kinase 2, probably acting as a signaling receptor of extracellular matrix component and required for cell expansion [72], was also up-regulated in AH cultivars. Remarkably, two probable plant transposases (Ptt/En/Spm families) (Oleur061Scf6180g00005.1, Oleur061Scf1200g04004.1) were up regulated in AH cultivars, suggesting an elevated transposon activity of these families of transposable element in AH roots. Finally, other genes related to membrane traffic, respiration, oxidoreductase activity, development or just unknown function were up-regulated in the roots of the AH group (Table 1).

From the set of 21 genes up-regulated in AL roots and down-regulated in the AH group, two ethylene (ET)-responsive TF, encoded by *ERF4* and *ERF34* genes (Oleur061Scf9036g01047.1, Oleur061Scf1024g00024.1), that binds to the GCC-box pathogenesis-related promoter element, and one ET-responsive element, the *ACO3* gene (Oleur061Scf1251g02004.1) coding for

an aminocyclopropane-1-carboxylate oxidase involved in the biosynthesis of the plant hormone ET, are worth to be highlighted. Additionally, in this set of genes a TF (*GAI1*) (Oleur061Scf5473g04020.1) that acts as a repressor of the GA signaling pathway was identified. The *SHR* (*SHORT-ROOT*) gene (Oleur061Scf3289g01014.1), that encodes another TF, was found to be up-regulated in AL roots as well. This gene regulates the radial organization of the root in *Arabidopsis* [73]. The *LBD30* gene (Oleur061Scf0088g04051.1), also up-regulated in AL cultivars, is involved in root development. In fact, overexpression of this gene produced shrunken root tips and disorganized columella cells in *Arabidopsis* [74]. Likewise, the *GSO1* gene (Oleur061Scf3112g01022.1) that codes a LRR receptor-like serine/threonine-protein kinase is involved in root growth. This gene regulates root growth through control of cell division and cell fate specification in *Arabidopsis* [75]. Finally, the *CSLC4* gene (Oleur061Scf2527g00016.1), coding for a xyloglucan glycosyltransferase 4 involved in synthesis of the non-cellulose component of the cell wall was also identified in this set of 21 up-regulated genes.

### 3.3. Effects of cultivar grouping in the distribution of the root microbiota

As mentioned above, the criterion to differentiate between AH and AL cultivars was the significant dominance of the genus *Actinophytocola* in the root endophytome of the AH group. This dominance was statistically proven when analyzing the  $\alpha$ -diversity of the endophytic bacterial community. Interestingly enough, the same (i.e. lower richness and diversity indices in the AH group than in AL cultivars) was also observed for the endophytic fungal community. However, no differences were found in the rhizosphere microbial communities (Table 3, Additional File 6: Fig. S4).

Surprisingly, when comparing the rhizosphere microbial (bacterial and fungal) composition by PERMANOVA and NMDS of the AH and AL cultivars (Table 3, Additional File 7: Fig. S5) both domains showed statistically different communities, an outcome *a priori* expected only for the endosphere. Therefore, the rhizosphere microbial communities of the AH and AL groups showed similar richness and  $\alpha$ -diversity values but with statistically distinguishable microbial assemblies ( $\beta$ -diversity).

### 3.4. The belowground microbiota of the AH and AL groups show different co-occurrence network topologies

Concerning the rhizosphere microbiota, similar co-occurrence network properties were obtained between AH and AL groups. The most remarkable difference was the much larger intra-modular connectivity (formerly known as ‘centralization of stress centrality’ and shown in Table 4 as CS) of the AH network (3.082) compared to that of the AL network (0.579). That is, the AH network had greater number, and more interconnected, module hubs (Table 4 and Fig. 2).

Regarding the root endosphere community two clearly different co-occurrence networks were obtained. Indeed, the AH network was more complex (higher avgCC and lower GD) and with much lower M value). As a whole, these indices were suggestive of a more interconnected network in the AH group. Additionally, the ‘percentage of positive edges’ (PEP) parameter was much lower in the AL group, indicating a higher proportion of negative interactions in this network (Table 4 and Fig. 2). Furthermore, an OTU belonging to *Actinophytocola* was a keystone node in the AH network, this OTU being the only module hub in the largest module of this network. In contrast, it was just a peripheral node in the AL network (Additional File 8: Fig. S6).

### 3.5. Connecting the olive root transcriptome with the endophytic microbial community

In light of the results so far presented, we investigated a potential correlation between DEGs and the different microbial profiles found in the root endosphere of AH and AL cultivars. Correlation analysis between the root endosphere microbial community and the set of DEGs previously identified in the AH and AL groups (8

FC,  $p$ -value  $\leq 0.01$ ; Tables 1 and 2) showed a strong ( $-0.6 \leq r \leq 0.6$ ) and significant ( $p < 0.05$ ) correlation with 111 OTUs. Two clearly different profiles could be observed (Fig. 3). On the one hand, a cluster with 82 OTUs belonging to nine different phyla, plus some unclassified bacteria and fungi, showed negative correlation with the up-regulated genes in the AH group (first row, yellow squares; Fig. 3) and positive correlation with the up-regulated genes in the AL group (first row, blue squares; Fig. 3). On the other hand, another cluster comprising 29 OTUs displayed the opposite correlation profile. Remarkably, 25 out of 29 OTUs in this cluster belonged to the phylum *Actinobacteria*, eight of them identified as *Actinophytocola* and eleven additional OTU unclassified at the genus level but assigned to the same family (i.e. *Pseudonocardiaceae*) as *Actinophytocola*.

The Pearson’s correlation analysis enabled to draw a first approximation to the interaction taking place between the root host transcriptome and the endophytic community. To go one step further in the search for keystone OTUs and genes in such interaction, a co-occurrence network analysis between the 60 DEGs and all OTUs observed in the root endosphere was performed. This analysis showed significant interactions between 32 DEGs (18 up-regulated and 14 down-regulated in the AH group, yellow and blue circles, respectively) and 19 OTUs (Fig. 4). Among all interactions observed, some of them were worth mentioning. For instance, OTU00021, belonging to *Actinophytocola*, showed a positive correlation with three up-regulated genes in the AH group (Oleur061Scf2041g01041.1, Oleur061Scf7501g00024.1 and Oleur061Scf3782g00023.1). The first one is related to response to biotic stress (functionally annotated as ‘Germin-like protein subfamily 1 member 16’) while the other two with unknown functions (annotated as ‘unknown protein’ and ‘Leishmanolysin-like peptidase’) (Fig. 4 and Table 1). The same OTU showed a negative correlation with two down-regulated genes in the AH group (Oleur061Scf1251g02004.1 and Oleur061Scf0724g05015.1, functionally annotated as ‘1-aminocyclopropane-1-carboxylate oxidase homolog’ and ‘Zinc finger, RING/FYVE/PHD-type’, respectively), the first one being key in the plant hormone ET biosynthesis pathway (Fig. 4 and Table 2). Furthermore, another OTU belonging to *Actinophytocola* (OTU00124) showed a positive correlation with two upregulated genes in the AH group (Oleur061Scf0191g00025.1 and Oleur061Scf0201g06012.1, both functionally annotated as ‘Thaumatococcus-like protein’). These two genes are related to response to biotic stress and antifungal activity (Fig. 4 and Table 1). The overall scenario shown in Fig. 4 is a negative correlation between all up-regulated genes in the AH group and all OTUs, except those ones belonging to the genus *Actinophytocola*. In contrast, the down-regulated genes in AH group showed just the opposite pattern.

## 4. Discussion

The interaction of the olive root transcriptome with the belowground microbiota has never been studied considering the holobiont context. Numerous studies have shown that the host

**Table 3**

Comparisons of  $\alpha$  and  $\beta$ -diversity indices between microbial communities of the *Actinophytocola* high-prevalence (AH) and low-prevalence (AL) groups.


	$\alpha$ -diversity				$\beta$ -diversity	
	Richness	Shannon	InvSimpson	Evenness	PERMANOVA	BETADISPER
Bacteria						
Endosphere	<b>2.46E-03</b>	<b>1.64E-02</b>	<b>2.35E-02</b>	<b>2.63E-02</b>	<b>1.00E-04</b>	<b>1.60E-02</b>
Rhizosphere	5.57E-01	4.61E-01	9.99E-01	5.12E-01	<b>1.00E-04</b>	6.44E-01
Fungi						
Endosphere	<b>4.24E-03</b>	<b>1.73E-03</b>	<b>1.39E-02</b>	1.27E-01	<b>5.10E-03</b>	4.69E-01
Rhizosphere	5.75E-01	3.05E-01	2.33E-01	2.85E-01	<b>1.00E-04</b>	<b>2.70E-02</b>

Mann-Whitney-Wilcoxon tests were performed to evaluate  $\alpha$ -diversity.  $\beta$ -diversity was compared using Bray-Curtis dissimilarities. Significant  $p$ -values are highlighted in bold type and italics.

**Table 4**  
The major topological properties of co-occurrence networks of *Actinophytocola* high-prevalence (AH) and low-prevalence (AL) groups.

Compartment	Group	Total nodes	Total links	Percentage of positive edges (PEP)	Similarity threshold (St)	R square of power-law	Average degree (avgK)	Average clustering coefficient (avgCC)	Average path distance (GD)	Modularity (M)	Centralization of stress centrality (CS)	Transitivity (Trans)
Endosphere	AH	129	491	97.56%	0.72	0.849	7.612	<b>0.305</b>	<b>2.890</b>	<b>12 (0.380)</b>	1.013	<b>0.286</b>
	AL	233	272	65.81%	0.83	0.903	2.335	<b>0.117</b>	<b>6.247</b>	<b>34 (0.834)</b>	0.848	<b>0.149</b>
Rhizosphere	AH	845	984	90.75%	0.85	0.904	2.329	<b>0.089</b>	<b>8.581</b>	120 (0.861)	<b>3.082</b>	<b>0.089</b>
	AL	758	948	91.56%	0.85	0.894	2.501	<b>0.099</b>	<b>8.722</b>	115 (0.857)	<b>0.579</b>	<b>0.150</b>

Figures in bold type and italics show statistically-significant differences between AH and AL groups in each compartment (root endosphere and rhizosphere) by Student's T-test.




**Fig. 2.** Co-occurrence networks of microbial communities of each compartment (root endosphere and rhizosphere) and each group (AH and AL). Each node corresponds to an OTU and were colored according to their phylum affiliation. Arrowheads represent module hubs and rhombus represent connectors. Lines represent positive (green) and negative (red) connections. AH: *Actinophytocola* high-prevalence group; AL: *Actinophytocola* low-prevalence group. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

genotype is decisive to explain the structure and composition of its resident microbial community [76,77]. However, a range of soil and environmental parameters, as well as diverse stresses, also affect the holobiont. Indeed, external factors influence the host genetic responses what in turn may determine decisive structural and functional changes in the associated microbiome [78,79]. Therefore, the global cross-talk established between the plant and its microbiome is not only crucial for the growth, development and fitness of the holobiont but also to efficiently cope with different (a)biotic constraints that must continuously face [15]. However, to our knowledge, this holistic approach has so far seldom accomplished, and available studies just focus either on the host side or on its associated microbiota [75–79]. In the case of olive trees, host whole-transcriptome responses have been analyzed under scenarios such abiotic [41,80] or biotic [46] stresses, or to unravel their effects over specific organs or tissues [45]. Concerning the olive microbiome, several studies have aimed to address questions such as the link between host tolerance/susceptibility to biotic stress and the structure and functioning of the residing

microbial community, or how the latter reacts against the presence of specific pathogens or insects [32,33,35]. One step further is to conduct dual or co-transcriptomic studies, but they mostly deal with a specific binomial as, for instance, the host transcriptome in response to an interaction with a given pathogen and vice versa [46,47]. However, studies in which the entire microbiota is evaluated in connection with the host global genetic response are lacking. Recently, metatranscriptomics (based on data mining of preexisting RNA-seq studies) has been implemented to offer a more global overview of the interaction between the indigenous microbial communities and the host genetic response, including the case of olive [48]. While these studies are providing relevant information, results might be biased and some relevant information overlooked because of limitations on the methodological approach implemented [49,81,82].

In the present study, we connect for the first time the olive root transcriptome to the belowground microbiota, aiming to identify host genes and microbial OTUs potentially crucial in the adaptation of the olive holobiont to a specific natural scenario. Our starting






**Fig. 3.** Heatmap with Pearson's correlations between the selected DEGs and the strongly and significantly correlated endophytic OTUs. Both OTUs and DEGs (see Tables 1 and 2 for functional annotation) were clustered with a complete-linkage clustering method and using Euclidean distance. bOtu's represent bacterial OTUs and fOtu's represent fungal OTUs. Strong correlation was considered when absolute value of Pearson correlation coefficient was  $\geq 0.6$ .

point was the available information on the structure and composition of the olive-associated belowground microbiota at the WOGC [20]. In this latter study, we identified *Actinophytocola* as the most prevalent bacterial genus among the surveyed cultivars, which has

been found linked to stress by water-supply deficiency. It must be emphasized that all cultivars in that study were grown under the same agronomical, pedological and climatic conditions, and that they underwent a prolonged period without water supply that



**Fig. 4.** Co-occurrence network showing the interactions between DEGs and endophytic OTUs. Only significant correlations were retained ( $-0.8 \leq r \leq 0.8$ ) according to the Random Matrix Theory (RMT) method (see co-occurrence network construction methodology). Circles represent up-regulated (yellow) and down-regulated (blue) genes in the AH group. Round squares and triangles represent bacterial and fungal OTUs, respectively, colored according to their deepest taxonomical annotation. Lines represent positive (green) and negative (red) connections. Gene-gene and OTU-OTU correlation lines were eliminated for clarification purpose. The main topological properties of the complete network were: Total nodes = 166, Total links = 551, PEP = 71.87%, St = 0.8, R2 of power-law = 0.861, avgK = 6.639, avgCC = 0.329, GD = 3.636, M = 17 (0.471), CS = 1.812, and Trans = 0.461. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

might be related to the high prevalence of *Actinophytocola* genus in the roots (Additional File 1: Table S1). However, the *Actinophytocola* relative abundance showed large variations among cultivars. Under these conditions, the genotype was the determinant factor explaining the differences found in the composition of the below-ground microbiota among cultivars. This led us to inquire whether cultivar-specific genetic responses (whole-transcriptome level) could be linked to the composition/structure of the microbial communities. Moreover, we also aimed to unravel differences in the network co-occurrence topologies of the outside (rhizosphere) and inside (root endosphere) communities of the two groups of cultivars selected. Finally, we investigated the feasibility to couple host gene expression profiles with the relative abundance of specific OTUs.

Our results demonstrated a correlation of the olive root microbiome with specific DEGs of the host. In fact, to our knowledge, this is the first study in which the interaction between specific host genes and precise root-inhabiting microbes is described. García and co-workers [81] carried out a similar study in the phyllosphere, although they used a different methodological approach (i.e. only transcriptomics data). Furthermore, a recent study used metatranscriptomics data to focus on the influence of the microbiome on differential gene expression in the tomato root endo-

sphere [49]. Unfortunately, however, this work did not provide details on which genes and microorganisms were interacting. Nevertheless, our results agreed with those of García and co-workers [81] in that the phylum *Actinobacteria* seems to play a key role when the plant holobiont is confronting a stressing situation. According to them, *Streptomyces* (phylum *Actinobacteria*) is particularly abundant in plants subjected to drought stress. Although this genus was present in the correlation analysis and positively correlated with a stress response gene (see OTU00023 and OTU00102 and gene *Oleur061Scf1329g05010.1* in Fig. 3), we did not find *Streptomyces* in the co-occurrence network (Fig. 4).

The roots of AH and AL cultivars displayed relevant differences in their global gene expression patterns, including genes related with plant defense response to stress. Strikingly, two Thaumatin-like coding genes, closely related with plant defense through apoplastic antifungal activity, showed higher expression in the AH group. Indeed, there was positive co-occurrence between these two genes and an *Actinophytocola* OTU (Fig. 4). This might be linked to a competitive relationship between fungi and bacteria, since the antifungal effect of thaumatin-like proteins might help *Actinophytocola* to flourish. Moreover, *Actinomycetales* are long known for their antagonistic behavior against root pathogenic fungi [83]. Nevertheless, it is very difficult to conclude whether the

abundance of *Actinophytocola* is cause or consequence of the observed higher expression of Thaumatin-like genes. Interestingly, a higher number of genes involved in cell wall synthesis or strength were also differentially expressed in AH roots. On the one hand, AH roots showed a higher transcriptional activity, as suggested not only by the higher expression of genes encoding proteins linked to protein turnover, but also by the upregulation of two Homeobox-leucine zipper TF. These TF play significant roles in the response to abiotic stress [71,84,85]. On the other hand, AL roots overexpressed several genes involved in root growth or development, sugar and water transport, lipid metabolism and pigment synthesis.

Regarding the root microbiome side, it is worth mentioning that a synchrony between endophytic bacteria and fungi was observed. Indeed, the opposite trend is frequently reported in the literature; that is, bacterial and fungal communities associated to the plant roots behave independently, at least in terms of  $\alpha$ -diversity, both in the rhizosphere [86] and the endosphere [87,88]. In fact, this lack of synchrony was previously observed in the microbial communities of the olive root endosphere when comparing two cultivars that were challenged with *V. dahliae* [35]. Interestingly, and in contrast with most of the studies, we found that the endophytic bacterial and fungal communities displayed lower richness and diversity in AH roots than in AL cultivars. This difference, at the bacterial level, was mainly due to the overwhelming dominance of the genus *Actinophytocola*. However, in the case of fungi, no significant difference in the evenness index between the two groups was observed (Table 3), which led to a decrease in diversity solely due to a reduction in the richness of fungal OTUs in AH varieties. Regarding the rhizosphere, no differences in richness and  $\alpha$ -diversity was found between the two groups of cultivars. Surprisingly, differences in  $\beta$ -diversity were observed in both the endosphere and the rhizosphere for the two microbial domains (Table 3, Additional File 7: Fig. S5). This result indicated that although the rhizosphere communities of both groups (AH and AL) were similar in richness and diversity, they showed significantly-different microbial compositions.

With regard to the interactions established within the microbial communities of each group of cultivars, it is worth mentioning the striking differences observed between network topologies of the endophytic communities (Fig. 2). The co-occurrence network of the AH group was more complex, with the individuals more interconnected (higher avgK and avgCC). In addition, these connections were stronger (lower GD and M) than in the AL network (Table 4). Considering the predominance of *Actinophytocola* in the AH group and the precedents on the adaptive capabilities of this genus to water stress conditions [50,51,53], the topological parameters found in this network suggested that the communities of each group responded to water stress (at least under the conditions found in the orchard at the time of sampling) in a very different way. In AH cultivars, the whole community seemed to work coordinately to adapt quickly to those stressing conditions. Accordingly to previous studies, higher and stronger connectivity (high avgK and avgCC linked to low GD) yielded networks with faster signal transmission among members of the community, and more rapid adaptation to any disturbance in the habitat [89–91]. Furthermore, this type of tightly-structured network (low GD and M) restricted the entry of new individuals with potential to alter it [92]. Therefore, such networks are generally more resistant and resilient to a/biotic stresses. In contrast, in the AL group a more heterogeneous topology was observed with larger distancing and compartmentalization of the community (high GD and M) [92,93]. We previously found this kind of network when the effect of the stressor was greater and the community aimed to “confine” it [35]. In fact, some microbial groups (modules in the network) responded in the opposite way to stress (thus a decrease in PEP was observed, Table 4).

Therefore, in this water stress scenario, results seemed to point to an increased competition among the members of the root endophytome in AL cultivars.

## 5. Conclusions

This study unveiled a significant interaction between specific genes up/down-regulated in olive roots and particular endophytes inhabiting this organ when trees were grown in the same agro-nomical and pedological scenario, and subjected to a period of water scarcity. Thus, a holistic view of the plant root-endophytic microbiome communication is here reported under specific environmental conditions. The roots of the two groups of olive cultivars examined showed significantly-different microbial profiles that strongly correlated with differential host gene expression patterns. The root transcriptome-microbiome “global communication” here described was solidly supported by both correlation and co-occurrence interaction analyses between host DEGs and microbial OTUs (Figs. 3 and 4), and showed a close coordination between a group of olive genes that negatively interacted with most members of its root endophytome. Moreover, these genes significantly decreased their expression in cultivars with the lowest relative abundance of the genus *Actinophytocola*, which seemed to play a key role. Our findings also suggested that the gene expression profile in roots of AL cultivars “facilitated” a more permissive endosphere environment for microbial colonization, thereby leading to more diverse and richer microbiome than the one found in the AH group. However, it also generated a more competitive environment (Table 4 and Fig. 2) for the members of this microbiome under the stressing conditions found at the time of sampling. In contrast, the AH group showed more restrictive conditions in its root endosphere what could have favored, among others, the flourishing of *Actinophytocola* under the observed conditions. Overall, the methodological approach here implemented can be useful to generate a specific “host transcriptome-microbiome signature” for any plant holobiont subjected to a particular environmental scenario. In future studies, the identification of specific olive root exudates and/or the evaluation of structural/biochemical modifications of the roots under specific stressing situations, among other approaches, will provide a more precise view on how the plant genotype affects its associated microbiome.

## Availability of data and materials

The datasets analyzed during the current study are available in the NCBI Sequence Read Archive (SRA) under the BioProject number PRJNA498945 (olive belowground microbiota), and in the Gene Expression Omnibus (GEO) repository, accession number GSE152236 (RNAseq).

## Funding

Supported by grants AGL2016-75729-C2-1-R and AGL2016-75729-C2-2-R from the Spanish Ministerio de Economía, Industria y Competitividad/Agencia Estatal de Investigación, and grant PID2019-106283RB-I00 from the Spanish Ministerio de Ciencia e Innovación/Agencia Estatal de Investigación, all of them co-financed by the European Regional Development Fund (ERDF). These funding sources had no involvement in this work preparation.

## Author's contributions

JMB, FL and MFL conceived and designed the study. AJFG, FL, and MFL implemented the methodology used. AJFG, JRT, PNB and



FL performed the bioinformatics analysis and analyzed the data. All authors contribute to the preparation of the original draft. AJFG, FL and JMB wrote, reviewed and edited the final version of the manuscript. Funding acquisition by FL, MFL and JMB. All authors read and approved the final manuscript.

### Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

### Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.csbj.2021.08.035>.

### References

- Loumou A, Giourga C. Olive groves: "The life and identity of the Mediterranean". *Agric Hum Values* 2003;20:87–95.
- Breton C, Guerin J, Ducatillon C, Médail F, Kull CA, Bervillé A. Taming the wild and "wilding" the tame: Tree breeding and dispersal in Australia and the Mediterranean. *Plant Sci* 2008;175(3):197–205.
- Besnard G, Terral JF, Cornille A. On the origins and domestication of the olive: a review and perspectives. *Ann Bot* 2018;121(3):385–403.
- Besnard G, Baradat P, Bervillé A. Genetic relationships in the olive (*Olea europaea* L.) reflect multilocal selection of cultivars. *Theor Appl Genet* 2001;102(2–3):251–8.
- FAO. FAOSTAT, Production Statistics. 2018. <http://faostat.fao.org> Accessed 22 Dec 2020.
- Ghanbari R, Anwar F, Alkharfy KM, Gilani AH, Saari N. Valuable nutrients and functional bioactives in different parts of olive (*Olea europaea* L.)—A review. *Int J Mol Sci* 2012;13(3):291–334.
- Gavahian M, Mousavi Khaneghah A, Lorenzo JM, Munekata PES, Garcia-Mantrana I, Collado MC, et al. Health benefits of olive oil and its components: impacts on gut microbiota antioxidant activities, and prevention of noncommunicable diseases. *Trends Food Sci Tech* 2019;88:220–7.
- Marcelino G, Hiane PA, Freitas KdeC, Santana LF, Pott A, Donadon JR, et al. Effects of olive oil and its minor components on cardiovascular diseases, inflammation, and gut microbiota. *Nutrients* 2019;11(8):1826. <https://doi.org/10.3390/nu11081826>.
- Bracci T, Busconi M, Fogher C, Sebastiani L. Molecular studies in olive (*Olea europaea* L.): Overview on DNA markers applications and recent advances in genome analysis. *Plant Cell Rep* 2011;30(4):449–62.
- Gucci R, Tattini M. Salinity tolerance in olive. *Hort Rev* 1997;21:177–214.
- Chartzoulakis KS. Salinity and olive: growth, salt tolerance, photosynthesis and yield. *Agri Water Manag* 2005;78(1–2):108–21.
- Sanzani SM, Schena L, Nigro F, Sergeeva V, Ippolito A, Salerno MG. Abiotic diseases of olive. *J Plant Pathol* 2012;94(3):469–91.
- Sebastiani L, Gucci R, Kerem Z, Fernández JE. Physiological responses to abiotic stresses. In: Rugini E, Baldoni L, Muleo R, Sebastiani L, editors. *The olive tree genome*. Springer International Publishing; 2016. p. 99–122.
- Mercado-Blanco J, Abrantes I, Barra Caracciolo A, Bevivino A, Ciancio A, Grenni P, et al. Belowground microbiota and the health of tree crops. *Front Microbiol* 2018;9.
- Trivedi P, Leach JE, Tringe SG, Sa T, Singh BK. Plant–microbiome interactions: from community assembly to plant health. *Nat Rev Microbiol* 2020;18(11):607–21.
- Pascale A, Proietti S, Pantelides IS, Stringlis IA. Modulation of the root microbiome by plant molecules: the basis for targeted disease suppression and plant growth promotion. *Front Plant Sci* 2020;10:1741.
- Jacoby RP, Koprivova A, Kopriva S. Pinpointing secondary metabolites that shape the composition and function of the plant microbiome. *J Exp Bot*. 2021; 72(1):57–69.
- Beckers B, Op De Beeck M, Weyens N, Boerjan W, Vangronsveld J. Structural variability and niche differentiation in the rhizosphere and endosphere bacterial microbiome of field-grown poplar trees. *Microbiome* 2017;5(1). <https://doi.org/10.1186/s40168-017-0241-2>.
- Berg C, Smalla K. Plant species and soil type cooperatively shape the structure and function of microbial communities in the rhizosphere. *FEMS Microbiol Ecol* 2009;68(1):1–13.
- Fernández-González AJ, Villadas PJ, Gómez-Lama Cabanás C, Valverde-Corredor A, Belaj A, Mercado-Blanco J, et al. Defining the root endosphere and rhizosphere microbiomes from the World Olive Germplasm Collection. *Sci Rep* 2019;9(1). <https://doi.org/10.1038/s41598-019-56977-9>.
- Garbeva P, van Elsas JD, van Veen JA. Rhizosphere microbial community and its response to plant species and soil history. *Plant Soil* 2008;302(1–2):19–32.
- Inceoglu Ö, Salles JF, van Elsas JD. Soil and cultivar type shape the bacterial community in the potato rhizosphere. *Soil Microbiol* 2012;63:460–70.
- Marschner P, Crowley D, Yang CH. Development of specific rhizosphere bacterial communities in relation to plant species, nutrition and soil type. *Plant Soil* 2004;261(1/2):199–208.
- Palla M, Turrini A, Cristani C, Caruso G, Avio L, Giovannetti M, et al. Native mycorrhizal communities of olive tree roots as affected by protective green cover and soil tillage. *Appl Soil Ecol* 2020;149:103520.
- Fausto C, Mininni AN, Sofo A, Crechchio C, Scagliola M, Dichio B, et al. Olive orchard microbiome: characterisation of bacterial communities in soil-plant compartments and their comparison between sustainable and conventional soil management systems. *Plant Ecol Div* 2018;11(5–6):597–610.
- Anguita-Maeso M, Trapero-Casas JL, Olivares-García C, Ruano-Rosa D, Palomo-Rios E, Jiménez-Díaz RM, et al. *Verticillium dahliae* inoculation and *in vitro* propagation modify the xylem microbiome and disease reaction to *Verticillium* wilt in a wild olive genotype. *Front Plant Sci* 2021;12:632689.
- Fernández-González AJ, Wentzien NM, Villadas PJ, Valverde-Corredor A, Lasa AV, Gómez-Lama Cabanás C, et al. Comparative study of neighboring Holm oak and olive trees belowground microbial communities subjected to different soil management. *PLoS ONE* 2020;15(8):e0236796.
- Miná D, Pereira JoséA, Lino-Neto T, Baptista P. Epiphytic and endophytic bacteria on olive tree phyllosphere: exploring tissue and cultivar effect. *Microb Ecol* 2020;80(1):145–57.
- Vergine M, Meyer JB, Cardinale M, Sabella E, Hartmann M, Cherubini P, et al. The *Xylella fastidiosa*-resistant olive cultivar "Leccino" has stable endophytic microbiota during the Olive Quick Decline Syndrome (OQDS). *Pathogens* 2020;9(1):35.
- Giampetrucci A, Baptista P, Morelli M, Cameirão C, Lino Neto T, Costa D, et al. Differences in the endophytic microbiome of olive cultivars infected by *Xylella fastidiosa* across seasons. *Pathogens* 2020;9(9):723.
- Anguita-Maeso M, Olivares-García C, Haro C, Imperial J, Navas-Cortés JA, Landa BB. Culture-dependent and culture-independent characterization of the olive xylem microbiota: effect of sap extraction methods. *Front Plant Sci* 2020;10:1708.
- Gomes T, Pereira JA, Lino-Neto T, Bennett AE, Baptista P. Bacterial disease induced changes in fungal communities of olive tree twigs depend on host genotype. *Sci Rep* 2019;9:5882.
- Abdelfattah A, Ruano-Rosa D, Cacciola SO, Li Destri Nicosia MG, Schena L, Papadopoulos NT. Impact of *Bactrocerca oleae* on the fungal microbiota of ripe olive drupes. *PLoS ONE* 2018;13(11):e0199403.
- Montes-Osuna N, Mercado-Blanco J. *Verticillium* wilt of olive and its control: what did we learn during the last decade? *Plants* 2020;9(6):735.
- Fernández-González AJ, Cardoni M, Gómez-Lama Cabanás C, Valverde-Corredor A, Villadas PJ, Fernández-López M, et al. Linking belowground microbial network changes to different tolerance level towards *Verticillium* wilt of olive. *Microbiome* 2020;8(1).
- Gros-Balthazard M, Besnard G, Sarah G, Holtz Y, Leclercq J, Santoni S, et al. Evolutionary transcriptomics reveals the origins of olives and the genomic changes associated with their domestication. *Plant J* 2019;100(1):143–57.
- Dastkar E, Soleimani A, Jafary H, de Dios Alche J, Bahari A, Zeinalbedini M, et al. Differential expression of genes in olive leaves and buds of ON- versus OFF-crop trees. *Sci Rep* 2020;10(1).
- Ramírez-Tejero JA, Jiménez-Ruiz J, Leyva-Pérez Mde la O, Barroso JB, Luque F. Gene expression pattern in olive tree organs (*Olea europaea* L.). *Genes* 2020;11(5):544.
- Bruno L, Picardi E, Pacenza M, Chiappetta A, Muto A, Gagliardi O, et al. Changes in gene expression and metabolic profile of drupes of *Olea europaea* L. cv Carolea in relation to maturation stage and cultivation area. *BMC Plant Biol* 2019;19(1). <https://doi.org/10.1186/s12870-019-1969-6>.
- Jiménez-Ruiz J, Leyva-Pérez Mde la O, Vidoy-Mercado I, Barceló A, Luque F. Transcriptomic time-series analysis of early development in olive from germinated embryos to juvenile tree. *BMC Genomics* 2018;19:824.
- Leyva-Pérez M.de la O, Valverde-Corredor A, Valderrama R, Jiménez-Ruiz J, Muñoz-Mérida A, Trelles O, et al. Early and delayed long term transcriptional changes and short-term transient responses during cold acclimation in olive leaves. *DNA Res* 2015;22(1):1–11. <https://doi.org/10.1093/dnares/dsu033>.
- Guerra D, Lamontanara A, Bagnaresi P, Orrù L, Rizza F, Zelasco S, et al. Transcriptome changes associated with cold acclimation in leaves of olive tree (*Olea europaea* L.). *Tree Genet Genomes* 2015;11(6):113.
- Bazakos C, Manioudaki ME, Sarropoulou E, Spano T, Kalaitzis P, Gerós H. 454 pyrosequencing of olive (*Olea europaea* L.) transcriptome in response to salinity. *PLoS ONE* 2015;10(11):e0143000.
- Giampetrucci A, Morelli M, Saponari M, Loconsole G, Chiumentoni M, Boscia D, et al. Transcriptome profiling of two olive cultivars in response to infection by the CoDiRO strain of *Xylella fastidiosa* subsp. *pauca*. *BMC Genomics* 2016;17(1). <https://doi.org/10.1186/s12864-016-2833-9>.
- Jiménez-Ruiz J, Leyva-Pérez M de la O, Schilirò E, Barroso JB, Bombarely A, Mueller L, et al. Transcriptomic analysis of *Olea europaea* L. roots during the *Verticillium dahliae* early infection process. *Plant Genome* 2017;10(1). <https://doi.org/10.3835/plantgenome2016.07.0060>.
- Leyva-Pérez M de la O, Jiménez-Ruiz J, Gómez-Lama Cabanás C, Valverde-Corredor A, Barroso JB, Luque F, et al. Tolerance of olive (*Olea europaea*) cv Frantoio to *Verticillium dahliae* relies on both basal and pathogen-induced differential transcriptomic responses. *New Phytol* 2018;217(2):671–86.
- Jiménez-Ruiz J, Leyva-Pérez M de la O, Gómez-Lama Cabanás C, Barroso JB, Luque F, Mercado-Blanco J. The transcriptome of *Verticillium dahliae* responds differentially depending on the disease susceptibility level of the olive (*Olea*



- europaea* L.) cultivar. *Genes* 2019;10(4):251. <https://doi.org/10.3390/genes10040251>.
- [48] Martí JM, Arias-Giraldo LF, Díaz-Villanueva W, Arnau V, Rodríguez-Franco A, Garay CP. Metatranscriptomic dynamics after *Verticillium dahliae* infection and root damage in *Olea europaea*. *BMC Plant Biol* 2020;20:79.
- [49] Chialva M, Ghignone S, Novero M, Hozzein WN, Lanfranco L, Bonfante P. Tomato RNA-seq data mining reveals the taxonomic and functional diversity of root-associated microbiota. *Microorganisms* 2020;8(1):38.
- [50] Ara I, Tsetseg B, Daram D, Suto M, Ando K. *Actinophytocola burenghanensis* sp. nov., isolated from Mongolian soil. *Int J Sys Evol Microbiol* 2016;66(7):2760–5.
- [51] Bouznada K, Bouras N, Schumann P, Spröer C, Sabaou N, Klenk HP. *Actinophytocola algeriensis* sp. nov., an actinobacterium isolated from Saharan soil. *Int J Sys Evol Microbiol* 2016;66(7):2760–5.
- [52] Cao C, Sun Y, Wu B, Zhao S, Yuan B, Qin S, et al. *Actinophytocola glycyrrhizae* sp. nov. isolated from the rhizosphere of *Glycyrrhiza inflata*. *Int J Sys Evol Microbiol* 2018; 68:2504–2508.
- [53] Sun HM, Zhang T, Yu LY, Lu XX, Mou XZ, Zhang YQ. *Actinophytocola gilvus* sp. nov., isolated from desert soil crusts, and emended description of the genus *Actinophytocola* Indananda et al. 2010. *Int J Syst Evol Microbiol* 2014; 64:3120–3125. DOI: 10.1099/ijss.0.061051-0
- [54] Indananda C, Matsumoto A, Inahashi Y, Takahashi Y, Duangmal K, Thamchaipenet A. *Actinophytocola oryzae* gen. nov., sp. nov., isolated from the roots of Thai glutinous rice plants, a new member of the family *Pseudonocardiaceae*. *Int J Syst Evol Microbiol* 2010;60:1141–6.
- [55] Ramírez-Tejero JA, Jiménez-Ruiz J, Serrano A, Belaj A, León L, de la Rosa R, et al. *Verticillium* wilt resistant and susceptible olive cultivars express a very different basal set of genes in roots. *BMC Genomics* 2021;22(1). <https://doi.org/10.1186/s12864-021-07545-x>.
- [56] Jiménez-Ruiz J, Ramírez-Tejero JA, Fernández-Pozo N, Leyva-Pérez Mde la O, Yan H, de la Rosa R, et al. Transposon activation is a major driver in the genome evolution of cultivated olive trees (*Olea europaea* L.). *Plant. Genome* 2020;13(1). <https://doi.org/10.1002/tpg2.v13.1.10.1002/tpg2.20010>.
- [57] Robinson MD, McCarthy DJ, Smyth GK. edgeR: a bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* 2010;26(1):139–40.
- [58] McMurdie PJ, Holmes S, Watson M. phyloseq: An R package for reproducible interactive analysis and graphics of microbiome census data. *PLoS ONE* 2013;8(4):e61217.
- [59] Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin P, O'Hara, et al. *Vegan*: community ecology package. R package *vegan*, vers. 2.2-1. 2015.
- [60] Wei T, Simko V. R package “corplot”: Visualization of a Correlation Matrix (Version 0.84). Available from <https://github.com/taiyun/corplot>. 2017.
- [61] Kolde R. pheatmap: Pretty heatmaps [Software] <https://cran.r-project.org/web/packages/pheatmap/index.html>. 2015.
- [62] Saeed AI, Sharov V, White J, Li J, Liang W, Bhagabati N, et al. TM4: a free, open-source system for microarray data management and analysis. *Biotechniques* 2003;34(2):374–8.
- [63] The Molecular Ecological Network Analyses Pipeline (MENAP) <http://ieg4.rccc.ou.edu/mena/main.cgi> [accessed Oct. 2020].
- [64] Deng Y, Jiang YH, Yang Y, He Z, Luo F, Zhou J. Molecular ecological network analysis. *BMC Bioinf* 2012;13:113.
- [65] Zhou J, Deng Y, Luo F, He Z, Tu Q, Zhi X, et al. Functional molecular ecological networks. *mBio* 2010;1(4). <https://doi.org/10.1128/mBio.00169-10>.
- [66] Zhou J, Deng Y, Luo F, He Z, Yang Y, Relman D. Phylogenetic molecular ecological network of soil microbial communities in response to elevated CO<sub>2</sub>. *mBio* 2011;2(4).
- [67] Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, et al. Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome Res* 2003;13:2498–504.
- [68] Serrano A, Rodríguez-Jurado D, López-Escudero FJ, Román B, Belaj A, de la Rosa R, et al. *Verticillium* wilt response of a olive core collection. The 7th international horticulture research conference, Northwest A&F University, China, July 2020, Fengwang Ma, Max Cheng, Eds. Yangling: Horticultural Research; 2020.
- [69] Moral J, Xaviér CJ, Viruega JR, Roca LF, Caballero J, Trapero A. Variability in susceptibility to anthracnose in the world collection of olive cultivars of Cordoba (Spain). *Front Plant Sci* 2017;8:1892.
- [70] Day RB, Shibuya N, Minami E. Identification and characterization of two new members of the *GRAS* gene family in rice responsive to N-acetylchitooligosaccharide elicitor. *Biochim Biophys Acta* 2003;1625(3):261–8.
- [71] Agalou A, Purwantomo S, Övernäs E, Johannesson H, Zhu X, Estiati A, et al. A genome-wide survey of HD-Zip genes in rice and analysis of drought-responsive family members. *Plant Mol Biol* 2008;66(1-2):87–103.
- [72] Wagner TA, Kohorn BD. Wall-associated kinases are expressed throughout plant development and are required for cell expansion. *Plant Cell* 2001;13(2):303–18.
- [73] Helariutta Y, Fukaki H, Wysocka-Diller J, Nakajima K, Jung J, Sena G, et al. The SHORT-ROOT gene controls radial patterning of the Arabidopsis root through radial signaling. *Cell* 2000;101(5):555–67.
- [74] Soyano T, Thitamadee S, Machida Y, Chua N-H. ASYMMETRIC LEAVES2-LIKE19/LATERAL ORGAN BOUNDARIES DOMAIN30 and ASL20/LBD18 regulate tracheary element differentiation in Arabidopsis. *Plant Cell*. 2008; 20:3359–3373.
- [75] Racolta A, Bryan AC, Tax FE. The receptor-like kinases GSO1 and GSO2 together regulate root growth in Arabidopsis through control of cell division and cell fate specification. *Dev Dyn* 2014;243(2):257–78.
- [76] Lundberg DS, Lebeis SL, Paredes SH, Yourstone S, Gehring J, Malfatti S, et al. Defining the core *Arabidopsis thaliana* root microbiome. *Nature* 2012;488(7409):86–90.
- [77] Bulgarelli D, Rott M, Schlaeppi K, Ver Loren van Themaat E, Ahmadinejad N, Assenza F, et al. Revealing structure and assembly cues for Arabidopsis root-inhabiting bacterial microbiota. *Nature* 2012;488(7409):91–5.
- [78] Delgado-Baquerizo M, Oliverio AM, Brewer TE, Benavent-González A, Eldridge DJ, Bardgett RD, et al. A global atlas of the dominant bacteria found in soil. *Science* 2018;359(6373):320–5.
- [79] Rath KM, Fierer N, Murphy DV, Rousk J. Linking bacterial community composition to soil salinity along environmental gradients. *ISME J* 2019;13(3):836–46.
- [80] Bazakos C, Manioudaki ME, Therios I, Voyiatzis D, Kafetzopoulos D, Awada T, et al. Comparative transcriptome analysis of two olive cultivars in response to NaCl-stress. *PLoS ONE* 2012;7(8):e42931.
- [81] García BJ, Labbé JL, Jones P, Abraham PE, Hodge I, Climer S, et al. Phytobiome and transcriptional adaptation of *Populus deltoides* to acute progressive drought and cyclic drought. *Phytobiomes Journal* 2018;2(4):249–60. <https://doi.org/10.1094/PBIOMES-04-18-0021-R>.
- [82] Dong C-J, Wang L-L, Li Q, Shang Q-M, Virolle M-J. Bacterial communities in the rhizosphere, phyllosphere and endosphere of tomato plants. *PLoS ONE* 2019;14(11):e0223847.
- [83] Lechevalier MP. Actinomycetes in agriculture and forestry. In: Goodfellow M, Williams ST, Mordarski M, editors. *Actinomycetes in biotechnology*. London: Academic Press; 1988. p. 327–58.
- [84] Gago GM, Almoguera C, Jordano J, Gonzalez DH, Chan RL. Habb-4, a homeobox-leucine zipper gene potentially involved in abscisic acid-dependent responses to water stress in sunflower. *Plant Cell Environ* 2002;25:633–40.
- [85] Wang Y, Henriksson E, Söderman E, Henriksson KN, Sundberg E, Engström P. The *Arabidopsis* homeobox gene, *ATHB16*, regulates leaf development and the sensitivity to photoperiod in *Arabidopsis*. *Dev Biol* 2003;264(1):228–39.
- [86] Wang Y, Xu X, Liu T, Wang H, Yang Y, Chen X, et al. Analysis of bacterial and fungal communities in continuous-cropping ramie (*Boehmeria nivea* L. Gaud) fields in different areas in China. *Sci Rep* 2020;10(1). <https://doi.org/10.1038/s41598-020-58608-0>.
- [87] Woźniak M, Grządziel J, Gałzka A, Frac M. Metagenomic analysis of bacterial and fungal community composition associated with *Paulownia elongata* × *Paulownia fortune*. *BioRes* 2019;14(4):8511–29. <https://doi.org/10.15376/biores.14.4.8511-8529>.
- [88] Kanasugi M, Sarkodee-Addo E, Ansong Omari R, Mohammad Golam Dastogeer K, Fujii Y, Oppong Abebrese S, et al. Exploring Rice Root Microbiome; the variation, specialization and interaction of bacteria and fungi in six tropic savanna regions in Ghana. *Sustainability* 2020;12(14):5835. <https://doi.org/10.3390/su12145835>.
- [89] Yang H, Li J, Xiao Y, Gu Y, Liu H, Liang Y, et al. An integrated insight into the relationship between soil microbial community and tobacco bacterial wilt disease. *Front Microbiol* 2017;8. <https://doi.org/10.3389/fmicb.2017.02179>.
- [90] Rybakova D, Mancinelli R, Wikström M, Birch-Jensen A-S, Postma J, Ehlers R-U, et al. The structure of the *Brassica napus* seed microbiome is cultivar-dependent and affects the interactions of symbionts and pathogens. *Microbiome* 2017;5(1). <https://doi.org/10.1186/s40168-017-0310-6>.
- [91] Jiemeng T, Meng D, Qin C, Liu X, Liang Y, Xiao Y, et al. Integrated network analysis reveals the importance of microbial interaction for maize growth. *Appl Microbiol Biotechnol* 2018;102:3805–18.
- [92] Cardinale M, Grube M, Erlicher A, Quehenberger J, Berg G. Bacterial networks and co-occurrence relationships in the lettuce root microbiota. *Environ Microbiol* 2015;17(1):239–52.
- [93] Delmas E, Besson M, Brice M-H, Burkle LA, Dalla Riva GV, Fortin M-J, et al. Analysing ecological networks of species interactions. *Biol Rev* 2019;94(1):16–36.