

# Hydrogen sulfide: A novel component in *Arabidopsis* peroxisomes which triggers catalase inhibition<sup>FA</sup>

Francisco J. Corpas<sup>1\*</sup>, Juan B. Barroso<sup>2</sup>, Salvador González-Gordo<sup>1</sup>, María A. Muñoz-Vargas<sup>1</sup> and José M. Palma<sup>1</sup>

1. Group of Antioxidants, Free Radicals and Nitric Oxide in Biotechnology, Food and Agriculture, Department of Biochemistry, Cell and Molecular Biology of Plants, Estación Experimental del Zaidín, CSIC, C/Profesor Albareda 1, E-18008 Granada, Spain

2. Group of Biochemistry and Cell Signaling in Nitric oxide, Department of Biochemistry and Molecular Biology, Campus “Las Lagunillas”, E-23071, University of Jaén, Jaén, Spain

\*Correspondence: Francisco J Corpas (javier.corpas@eez.csic.es)

doi: 10.1111/jipb.12779

**Abstract** Plant peroxisomes have the capacity to generate different reactive oxygen and nitrogen species (ROS and RNS), such as H<sub>2</sub>O<sub>2</sub>, superoxide radical (O<sub>2</sub><sup>•-</sup>), nitric oxide and peroxynitrite (ONOO<sup>-</sup>). These organelles have an active nitro-oxidative metabolism which can be exacerbated by adverse stress conditions. Hydrogen sulfide (H<sub>2</sub>S) is a new signaling gasotransmitter which can mediate the posttranslational modification (PTM) persulfidation. We used *Arabidopsis thaliana* transgenic seedlings expressing cyan fluorescent protein (CFP) fused to a canonical peroxisome targeting signal 1 (PTS1) to visualize peroxisomes in living cells, as well as a specific fluorescent probe which showed that peroxisomes contain H<sub>2</sub>S. H<sub>2</sub>S was also detected in chloroplasts under glyphosate-induced oxidative stress conditions. Peroxisomal enzyme activities, including catalase,

photorespiratory H<sub>2</sub>O<sub>2</sub>-generating glycolate oxidase (GOX) and hydroxypyruvate reductase (HPR), were assayed *in vitro* with a H<sub>2</sub>S donor. In line with the persulfidation of this enzyme, catalase activity declined significantly in the presence of the H<sub>2</sub>S donor. To corroborate the inhibitory effect of H<sub>2</sub>S on catalase activity, we also assayed pure catalase from bovine liver and pepper fruit-enriched samples, in which catalase activity was inhibited. Taken together, these data provide evidence of the presence of H<sub>2</sub>S in plant peroxisomes which appears to regulate catalase activity and, consequently, the peroxisomal H<sub>2</sub>O<sub>2</sub> metabolism.

**Edited by:** Jianping Hu, Michigan State University, USA

**Received** Sept. 27, 2018; **Accepted** Jan. 14, 2019; **Online on** Jan. 16, 2019

FA: Free Access

Research Article

Free Access

## INTRODUCTION

Multifunctional hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) and nitric oxide (NO) molecules are involved in physiological processes, as well as responses to adverse environmental conditions (Neill et al. 2002; Corpas 2015; del Río 2015; da Silva et al. 2017; Corpas and Palma 2018). Though characterized by a very simple chemical structure, both these molecules are capable of generating reactive oxygen and nitrogen species (ROS and RNS).

Hydrogen sulfide (H<sub>2</sub>S), the simplest thiol found in animal and plant cells, has been known to be toxic for some time. However, it has recently been found to have properties similar to those of multifunctional signaling molecules (NO and H<sub>2</sub>O<sub>2</sub>) (Lisjak et al. 2013; Gotor et al.

2015; Hancock and Whiteman 2016; Li et al. 2016; Yamasaki and Cohen 2016; Filipovic and Jovanovic 2017). As with NO and H<sub>2</sub>O<sub>2</sub>, exogenous applications of H<sub>2</sub>S in plants have been shown to counteract the toxic effects of stresses, such as heavy metal and salinity (Zhang et al. 2010; Chen et al. 2014, 2015, 2017; Ali et al. 2014; Bharwana et al. 2014; Kharbech et al. 2017; Corpas et al. 2019). In plant systems, several enzymes are capable of generating H<sub>2</sub>S, which is part of the cysteine (Cys) metabolism. These enzymes, including L- and D-cysteine desulfhydrase (L-DES/D-DES), sulfite reductase (SiR), cyano alanine synthase (CAS) and cysteine synthase (CS) (Li et al. 2013; Calderwood and Kopriva 2014; Hancock and Whiteman 2014), are present in different subcellular compartments, such as cytosol, chloroplasts

and mitochondria (Gotor et al. 2015; Hancock and Whiteman 2016). For example, endogenous H<sub>2</sub>S content has recently been reported to increase during sweet pepper fruit ripening, which is associated with an increase in cytosolic L-cysteine desulphydrase (L-DES) activity (Muñoz-Vargas et al. 2018).

With their active nitro-oxidative metabolisms, plant peroxisome organelles are necessary in multiple biochemical pathways in all phases of plant development, from seed germination to plant senescence and also in response to adverse environmental conditions (del Río et al., 2002; Hu et al. 2012; Corpas et al. 2017; Kao et al. 2018; Palma et al. 2018). To our knowledge, no information exists on the presence of H<sub>2</sub>S in plant or animal peroxisomes. However, proteomic studies of animals and plants have identified catalase as a potential target of persulfidation (Mustafa et al. 2009; Aroca et al. 2015), which is a post-translational modification (PTM) mediated by H<sub>2</sub>S that modulates the function of target proteins (Iciek et al. 2015; Ju et al. 2017). Thus, this study mainly aims to evaluate the presence of H<sub>2</sub>S in plant peroxisomes and its potential biochemical implications.

## RESULTS


In 2014, Peng et al. devised a series of pyridine disulfide-based fluorescent Washington State Probes (WSP1 to WSP5) to detect H<sub>2</sub>S in both aqueous solutions and cell images. H<sub>2</sub>S can undergo dual nucleophilicity which facilitates a tandem nucleophilic substitution-cyclization reaction, enables fluorophore release and activates fluorescence. These WSPs also exhibited greater sensitivity and selectivity in relation to H<sub>2</sub>S as compared to other cellular sulfur species, such as cysteine (Cys), and reduced glutathione (GSH). Using fluorescence microscopy, Peng et al. (2014) observed the presence of H<sub>2</sub>S in human HeLa living cells pre-incubated with these WSP fluorescence probes.

Using confocal laser scanning microscope (CLSM) to analyze 10-d-old *Arabidopsis thaliana* transgenic seedlings expressing the cyan fluorescent protein (CFP) fused to canonical peroxisome targeting signal 1 (PTS1), we observed peroxisomes in the form of spherical spots (green color) distributed randomly throughout *Arabidopsis* root tip cells (Figure 1A). We used the WSP-5 fluorescent probe to specifically detect cellular

H<sub>2</sub>S (Peng et al. 2014; Yu et al. 2014). Using this probe and CLSM with the CFP-PTS1 *Arabidopsis* transgenic line, we observed intense red fluorescence corresponding to H<sub>2</sub>S in spherical spots on the root tip cells (Figure 1B), with a pattern analogous to that of CFP-PTS1 (Figure 1A). Figure 1C shows a merged image of panels A and B, with a significant correspondence of both punctate distributions, indicating that H<sub>2</sub>S is present in *Arabidopsis* peroxisomes. A similar analysis was carried out on the guard cells of green *Arabidopsis* cotyledons (Figure 1D, G), with chlorophyll autofluorescence also enabling us to observe the presence of chloroplasts (Figure 1F). Figure 1G shows the merged image of panels D to F, which indicate that H<sub>2</sub>S is located in peroxisomes and the cytosol.

Although CFP excitation/emission wavelengths do not theoretically overlap in WSP-5 when used to detect H<sub>2</sub>S, this was evaluated at the experimental level in order to rule out this possibility. 10-d-old *Arabidopsis* seedlings expressing CFP-PTS1 were, therefore, used without pre-incubation with fluorescent probe WSP-5 to detect H<sub>2</sub>S. Thus, Figure S1A shows the detection by CLSM of peroxisomes in the root tip of an *Arabidopsis* seedling expressing CFP-PTS1 using excitation and emission wavelengths of 459 and 475 nm, respectively. On the other hand, Figure S1B shows the same *Arabidopsis* root area with excitation and emission wavelengths set at 502 nm and 525 nm, respectively. Under these conditions, no fluorescent signal was observed, which corroborated the absence of overlap between CFP and WSP-5.

In order to evaluate the specificity and efficiency of WSP-5 in the detection of H<sub>2</sub>S in *Arabidopsis* seedlings, additional controls using *Arabidopsis* wild-type (WT) seedlings were carried out. Figure 2A–D shows a primary root of 10-d-old *Arabidopsis* seedlings incubated with increasing concentrations (0.0, 0.1, 0.5 and 2 mmol/L) of sodium hydrosulfide NaHS which is a recognized H<sub>2</sub>S donor molecule in animal and plant systems (Wang et al. 2010; Aroca et al. 2015; Yang et al. 2016; Kharbech et al. 2017; Zhang et al. 2017; Muñoz-Vargas et al. 2018). Thus, an increase in the red fluorescence signal corresponding to the presence of H<sub>2</sub>S was also observed. Furthermore, when *Arabidopsis* seedlings were pre-incubated with 0.1 mmol/L NaHS and 0.1 mmol/L hypotaurine (a H<sub>2</sub>S scavenger), the fluorescence attributable to H<sub>2</sub>S decreased significantly in the root (Figure 2E) as compared to the seedling treated




**Figure 1.** Representative images illustrating the CLSM *in vivo* detection of H<sub>2</sub>S (red color) and peroxisomes (green color) in root tips (A to C) and guard cells (D to G) of 10-d-old *Arabidopsis* seedlings expressing CFP-PTS1 (A) and (D), Fluorescence punctates (green) attributable to CFP-PTS1, indicating the localization of peroxisomes. (B) and (E), Fluorescence punctates (red) attributable to H<sub>2</sub>S detection in the same area. (C) Merged image of (A) and (B) showing colocalized fluorescence punctates (yellow). (F) Chlorophyll autofluorescence (blue) demonstrating location of chloroplasts. (G) Merged images of (D) to (F). H<sub>2</sub>S (red color) was detected by using 5 μM WSP-5. Arrows indicate representative punctuate spots corresponding to the colocalization of H<sub>2</sub>S with peroxisomes. Asterisks indicate localization of H<sub>2</sub>S in the cytosol.

with 0.1 mmol/L NaHS alone (Figure 2B). Figure 2F shows the change in the fluorescence intensity of H<sub>2</sub>S (red color) in the primary roots of *Arabidopsis* seedlings (Figure 2A–E). These data confirm that WSP-5 is a reliable tool for analyzing H<sub>2</sub>S in *Arabidopsis* seedlings.

To assess how H<sub>2</sub>S content and cellular distribution are affected under stress conditions, *Arabidopsis* seedlings were grown in the presence of glyphosate, a herbicide which triggers oxidative stress in peroxisomal metabolism (de Freitas-Silva et al. 2017). Figure 3 shows that H<sub>2</sub>S detected in the green cotyledons (Figure 3A–H) and roots (Figure 3I–N) of 14-d-old *Arabidopsis* seedlings grown under optimal conditions or in the presence of 20 μM glyphosate. Under optimal conditions, as previously described (Figure 1), H<sub>2</sub>S was mainly detected in peroxisomes and the cytosol. However, under glyphosate-induced oxidative stress, red fluorescence corresponding to H<sub>2</sub>S content increased significantly in green cotyledons and roots

(Figure 3D, L). In addition to its presence in green cotyledons and peroxisomes, H<sub>2</sub>S was also detected in other rounded structures corresponding to chloroplasts (Figure 3D, F, H).

We used catalase to evaluate the potential impact of H<sub>2</sub>S on the peroxisomal metabolism. Proteomic analysis has identified catalase, which is the main peroxisomal antioxidant enzyme, as a target of persulfidation in mouse liver (Mustafa et al. 2009) and *Arabidopsis* leaves (Aroca et al. 2015, 2017). Therefore, we analyzed enzymatic catalase activity under *in vitro* conditions using a NaHS concentration gradient. Figure 4A shows the impact of H<sub>2</sub>S inhibition on catalase activity, with doses of NaHS ranging from 0.1 to more than 4 mmol/L, which led to reductions of 29% and 88% in catalase activity, respectively. Accordingly, the H<sub>2</sub>S donor inhibited catalase activity by 50% (IC<sub>50</sub>) at a concentration of approximately 0.9 mmol/L. We also studied photorespiratory H<sub>2</sub>O<sub>2</sub>-generating glycolate oxidase (GOX), another peroxisomal enzyme



**Figure 2.** Representative images illustrating the CLSM *in vivo* detection of H<sub>2</sub>S (red color) in primary roots of 10-d-old *Arabidopsis thaliana* wild-type seedlings pre-incubated with different concentrations of NaHS (0.0, 0.1, 0.5 and 2 mmol/L) (A to D, respectively) and pre-incubated with 0.1 mmol/L NaHS plus 0.1 mmol/L hypotaurine (HyT), a H<sub>2</sub>S scavenger (E)


The fluorescence intensity of the red WSP-5 signal in roots of panels (A) to (E) was determined using Image J software and is expressed as arbitrary units (A.U.) (F).

(Oikawa et al. 2015; Hagemann and Bauwe 2016), whose activity decreased by 22% at a concentration of 4 mmol/L NaHS (Figure 4B). In addition, hydroxypyruvate reductase (HPR) activity, which is also involved in the photo-respiratory pathway, was barely affected by these concentrations (Figure 4C). To corroborate the inhibitory effect of H<sub>2</sub>S on *Arabidopsis* catalase activity, catalase from green pepper fruit-enriched samples and pure catalase from bovine liver were also assayed (Figure 5A, B, respectively). In both cases, catalase activity was inhibited in the presence of NaHS, with an IC<sub>50</sub> of 0.51 mmol/L for the pepper fruit catalase and 0.98 mmol/L for bovine liver catalase.

## DISCUSSION

Persulfidation is a post-translational modification (PTM) in which the thiol group (-SH) of a specific cysteine (Cys) residue is converted into a persulfide group (-SSH). Thus, protein persulfidation is an oxidative PTM that can

mediate signaling events caused by H<sub>2</sub>S (Filipovic and Jovanovic 2017; Filipovic et al. 2018; Corpas et al. 2019). Using the modified biotin switch method, 106 putative target proteins, which can be persulfidated in Cys residues in 30-d-old *Arabidopsis thaliana* leaves, were initially identified (Aroca et al. 2015). *In vitro* assays in the presence of the H<sub>2</sub>S donor NaHS in selected proteins showed inactivated glutamine synthetase (GS) activity, while ascorbate peroxidase (APX) and glyceraldehyde 3-phosphate dehydrogenase (GAPDH) activity increased significantly (Aroca et al. 2015). Using an improved tag switch method with a biotin-linked cyanoacetate (CN-biotin) to form stable thiol-ether conjugates (Zhang et al. 2014) combined with LC-MS/MS analysis, Aroca et al. (2017) found that the number of putative persulfidated proteins in *Arabidopsis* leaves increased to 2,015. After inspecting the reported list of *Arabidopsis* persulfidated proteins, 17 peroxisomal proteins, representing 0.8% of the total, were identified, including three catalase isozymes, several fatty acid  $\beta$ -oxidation enzymes and a few enzymes involved in photorespiration pathways (see



**Figure 3. Representative images illustrating the CLSM *in vivo* detection of H<sub>2</sub>S (red) in green cotyledons (panels A to H) and roots (panels I to N) of 14-d-old *Arabidopsis* seedlings expressing CFP-PTS1 (green) grown in the presence of 20 μM glyphosate**

(A), (B), (I) and (J), Fluorescence punctates (green) attributable to CFP-PTS1, indicating the localization of peroxisomes. (C), (D), (K) and (L), Red fluorescence attributable to H<sub>2</sub>S detection in the same area. (E) and (F), Chlorophyll autofluorescence (blue) demonstrating location of chloroplasts. (G), (H), (M) and (N) merged images. (O), fluorescence intensity of the red WSP-5 signal in green cotyledons (C and D) and in roots (K and L)


**Table 1).** However, the effect of persulfidation was not analyzed in any of these proteins. We therefore evaluated the possible presence of H<sub>2</sub>S in *Arabidopsis* peroxisomes using cell imaging techniques and, with the aid of *in vitro* assays, we also studied the potential impact of H<sub>2</sub>S on some of the peroxisomal proteins identified, including antioxidant catalase, photorespiratory H<sub>2</sub>O<sub>2</sub>-generating GOX and hydroxypyruvate reductase (HPR).

### H<sub>2</sub>S is present in plant cell cytosols, peroxisomes and chloroplasts

In previous studies, using *Arabidopsis* seedlings expressing CFP-PTS1 and specific fluorescent probes, we detected the presence of various molecules involved in the peroxisomal metabolism of ROS and RNS, such as superoxide radicals (O<sub>2</sub><sup>-</sup>), peroxynitrite (ONOO<sup>-</sup>) and nitric oxide (NO), as well as the presence of calcium which is essential for NO generation (Corpas and Barroso 2014, 2018; Corpas et al. 2009). On the other hand, WSP fluorescent probes have proven useful to visualize H<sub>2</sub>S in animal HeLa cells (Peng et al. 2014) and tomato roots (Li et al. 2014). We used fluorescent WSP-5, which has a faster fluorescence turn-on rate and more sensitive detection limits as compared to other WSPs, to study the potential presence of H<sub>2</sub>S in plant peroxisomes. The data obtained, together with those from controls (Figure 2), indicate that the WSP-5 fluorescent probe is useful for detecting H<sub>2</sub>S in plant systems. Similar observations have been made in relation to tomato roots using fluorescent probe WSP-1 (Li et al. 2014). In addition, using *Arabidopsis* seedlings expressing CFP-PTS1, which enable peroxisomes to be visualized in living cells incubated with fluorescent probe WSP-5, as well as CLSM, we observed a red fluorescent signal attributable to H<sub>2</sub>S which was present in the cytosol and peroxisomes of green cotyledons and roots of *Arabidopsis* seedlings grown under optimal conditions. The presence of H<sub>2</sub>S in other cell compartments cannot be ruled out, as several enzymatic sources of H<sub>2</sub>S present in cytosols, chloroplasts and mitochondria are involved in the cysteine metabolism (Papenbrock et al. 2007; Alvarez et al. 2010;

Birke et al. 2015; Li et al. 2016), although none appear to be located in peroxisomes. To explore this possibility, *Arabidopsis* seedlings were grown in the presence of glyphosate which affects the shikimate pathway, reduces aromatic amino acid synthesis and disturbs plant growth and secondary metabolites which compromises auxin and salicylate biosynthesis occurs (Tzin and Galili 2010; Peek and Christendat 2015). We previously demonstrated that, in *Arabidopsis* seedlings, glyphosate triggers oxidative stress in the peroxisomal metabolism and in the oxidative phase of the pentose phosphate pathway (OxPPP) (de Freitas-Silva et al. 2017). Under these oxidative conditions, CLSM showed that H<sub>2</sub>S content increased in both green cotyledons and roots. However, interestingly, its location in chloroplasts is very much in line with that of H<sub>2</sub>S-generating sulfite reductase (Khan et al. 2010). On the other hand, a recent study identified a novel pathway that produces H<sub>2</sub>S from D-cysteine in animals, involving D-amino acid oxidase (DAO) and 3-mercaptopyruvate sulfurtransferase (3MST), which are present in peroxisomes and mitochondria, respectively (Kimura 2015). H<sub>2</sub>S could be present in other subcellular compartments of *Arabidopsis* due to the chemical properties of this highly lipophilic molecule which easily spreads throughout the lipid bilayer of cell membranes (Mathai et al. 2009; Cuevasanta et al. 2017). Thus, given an estimated H<sub>2</sub>S mobility of 0.5 cm/s in the lipid bilayer, or around four orders of magnitude faster than water (Mathai et al. 2009; Riahi and Rowley 2014), H<sub>2</sub>S accumulated in *Arabidopsis* peroxisomes could originate from cytosol and chloroplasts. It has not been determined whether peroxisomal H<sub>2</sub>S is endogenously generated or imported, although both are possible. It is worth noting that plant peroxisomes contain enzymatic and non-enzymatic components such as glutathione reductase (GR), sulfite oxidase (SO), glutathione (GSH), S-nitrosoglutathione (GSNO) and sulfite (Jiménez et al. 1997; Hänsch and Mendel 2005; Hänsch et al. 2006; Corpas and Barroso 2015), which are involved in the sulfur metabolism.

determined using Image J software and is expressed as arbitrary units (A.U). H<sub>2</sub>S (red color) was detected by using 5 μM WSP-5. Arrows indicate representative punctuate spots corresponding to the colocalization of H<sub>2</sub>S with peroxisomes. Squares with broken lines indicate localization of H<sub>2</sub>S in the chloroplasts.



**Figure 4. Effect of NaHS (H<sub>2</sub>S donors) on several peroxisomal enzymes of *Arabidopsis* seedlings**

Peroxisomal enzymatic activity assays for (A) catalase expressed as μmol H<sub>2</sub>O<sub>2</sub> · min<sup>-1</sup> · mg<sup>-1</sup> protein, (B) glycolate oxidase (GOX) expressed as μmol Glyoxylate phenylhydrazone · min<sup>-1</sup> · mg<sup>-1</sup> protein, and (C) hydroxypyruvate reductase (HPR) expressed as nmol NADH · min<sup>-1</sup> · mg<sup>-1</sup> protein. Data are means ± SE of at least three replicates. Differences from control values were significant at *P* < 0.05.


### Inhibition of catalase by H<sub>2</sub>S

The presence of H<sub>2</sub>S in specific organelles is also evidenced by the identification of the protein post-translational modification persulfidation (also known as S-sulfhydration) which affects susceptible reactive

cysteine residues and converts Cys-SH groups into Cys-SSH groups (Filipovic 2015; Paul and Snyder 2015; Filipovic et al. 2018). Using a modified biotin switch method combined with liquid chromatography-tandem mass spectrometry, 39 and 2,015 endogenously persulfidated proteins have been identified in mouse liver (Mustafa et al. 2009) and *A. thaliana* leaves (Aroca et al. 2015, 2017), respectively. In both these studies, catalase, which is exclusively located in peroxisomes, was observed to be a target for persulfidation, a finding that corroborates the functional presence of H<sub>2</sub>S in these organelles. However, to our knowledge, no information exists on the specific effects of persulfidation on catalase activity. It is important to point out that we used high concentrations of the H<sub>2</sub>S donor, as NaHS, which is immediately hydrolyzed in aqueous solution, establishes an equilibrium between H<sub>2</sub>S, HS<sup>-</sup> and S<sub>2</sub><sup>-</sup> species. Moreover, after this equilibrium is established, H<sub>2</sub>S is volatilized, which reduces the concentration of sulfur species in solution. Air oxidation of HS<sup>-</sup> catalyzed by the presence of trace metals in aqueous solution also reduces the actual concentration of H<sub>2</sub>S in solution (Hughes et al. 2009). However, as mentioned above, NaHS is the most commonly used H<sub>2</sub>S donor in animal and plant systems.

Catalase, which regulates peroxisomal H<sub>2</sub>O<sub>2</sub> content, is exclusively located in peroxisomes; the inhibition of catalase by persulfidation is in line with previous studies which indicate that catalase is also targeted by other PTMs mediated by catalase-inhibiting NO-derived molecules, such as S-nitrosylation and nitration (Clark et al. 2000; Begara-Morales et al. 2013; Chaki et al. 2015; Hu et al. 2015; Titov and Osipov 2017). This suggests that catalase is strictly regulated by H<sub>2</sub>S and NO, which is particularly important under adverse conditions given the nitro-oxidative environment inside peroxisomes (Corpas et al. 2017). The other peroxisomal enzymes assayed under similar *in vitro* conditions behaved differently, with GOX experiencing an inhibition of 22% at higher concentrations of NaHS, while HPR was virtually unaffected, indicating that the inhibitory effect on catalase is specific.

It is possible to conclude that, although plant peroxisomes are known to contain biologically active ROS and RNS metabolisms, the presence of H<sub>2</sub>S opens up new questions about their role under physiological and stress conditions. Catalase, an essential enzyme in living organisms and one of the first peroxisomal



**Figure 5. Effect of NaHS (H<sub>2</sub>S donors) on catalase activity from different origins**

(A) 0–50% enriched (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> protein fraction obtained from green pepper fruits. (B) Commercial pure catalase from bovine liver. Catalase activity is expressed as μmol H<sub>2</sub>O<sub>2</sub> · min<sup>-1</sup> · mg<sup>-1</sup> protein. Data are means ± SE of at least four replicates. \*Differences from control values were significant at  $P < 0.05$ .

**Table 1. List of peroxisomal persulfidated proteins found in leaves of *Arabidopsis thaliana* (Aroca et al. 2015, 2017) with its corresponding Uniprot accession number**

Peroxisomal protein	Uniprot accession number
Catalase 1	Q96528
Catalase 2	P25819
Catalase 3	Q42547
Glycerate dehydrogenase HPR, peroxisomal	Q9C9W5
Glycolate oxidase 1	Q9LRR9
3-ketoacyl-CoA thiolase 1	Q8LF48
3-ketoacyl-CoA thiolase 2	Q56WD9
3-ketoacyl-CoA thiolase 5	Q570C8
Peroxisomal fatty acid β-oxidation multifunctional protein AIM1	Q9ZPI6
Peroxisomal acyl-coenzyme A oxidase 1	O65202
Acetate/butyrate–CoA ligase AAE7	Q8VZF1
NADP-isocitrate dehydrogenase	Q9SLK0
Fatty acid β-oxidation multifunctional protein MFP2	Q9ZPI5
Acyl-coenzyme A oxidase 3	PoCZ23
Acyl-coenzyme A oxidase 4	Q96329
β-glucosidase 26	O64883
Enoyl-CoA hydratase 2	Q8VYI3

antioxidant enzymes to be characterized, catalyzes the decomposition of hydrogen peroxide ( $2\text{H}_2\text{O}_2 \rightarrow 2\text{H}_2\text{O} + \text{O}_2$ ) (Mhamdi et al. 2012; Corpas 2015; Su et al. 2018). The presence of H<sub>2</sub>S, which regulates catalase activity, supports the hypothesis that plant peroxisomes contain an active metabolism of reactive sulfur species (RSS) (Corpas and Barroso 2015). Previous studies have reported the presence of other enzymes, such as glutathione reductase (GR) and sulfite oxidase (SO), involved in the sulfur metabolism (Nowak et al. 2004; Hänsch et al. 2006, 2007). SO, which catalyzes the conversion of sulfite to sulfate with a concomitant generation of H<sub>2</sub>O<sub>2</sub>, appears to protect catalase, which is inhibited by low concentrations of sulfite (Veljović-Jovanović et al. 1998; Hänsch et al. 2007). In summary, our findings provide new evidence of the complexity of the peroxisomal metabolism in plants, in which H<sub>2</sub>S can be regarded as a new regulatory molecule that may be involved in crosstalk between peroxisomes and other subcellular compartments, especially under nitro-oxidative stress conditions. These data could also corroborate the potential presence of H<sub>2</sub>S in animal peroxisomes, as animal catalase is also reported to be persulfidated (Mustafa et al. 2009) and, consequently, inhibited, as shown in this study (Figure 4B). Thus, this situation could be similar to that in relation to peroxisomal superoxide dismutase (SOD) which was



first described in plant peroxisomes, a discovery which was not questioned until CuZn-SOD was detected in animal peroxisomes (Corpas et al. 2017).

## MATERIALS AND METHODS

### Arabidopsis growth conditions

Wild-type and transgenic seeds of *Arabidopsis thaliana* expressing cyan fluorescent protein (CFP) fused to a canonical peroxisome targeting signal 1 (PTS1) (Nelson et al. 2007) were surface-sterilized for 5 min using a solution of 70% ethanol containing 0.1% SDS. Then, the seeds were kept in sterile water containing 20% bleach and 0.1% SDS for 20 min and washed several times in sterile water. The seeds were sown for 2 d at 4°C in the dark on Petri plates containing 4.32 g/L Murashige and Skoog basal medium (Sigma), 1% sucrose and 0.8% phytoagar, with a pH of 5.5 (Corpas and Barroso 2014b). The *Arabidopsis* seeds were then grown for 10 d at 16 h light, 22°C/8 h dark, at 18°C (long day conditions) under a light intensity of 100 µE/m<sup>2</sup>/s (Corpas and Barroso 2017). California-type green sweet pepper (*Capsicum annuum* L.) fruits were provided for Syngenta Seeds Ltd. (El Ejido, Almería, Spain). Pure catalase from bovine liver was a product of Sigma. For the experiments with glyphosate stress, seeds were grown directly on MS medium plates with and without 20 µM glyphosate for 14 d under long-day conditions (de Freitas-Silva et al. 2017).

### Crude extracts from plant samples

*Arabidopsis* seedlings were collected, pooled and frozen in liquid nitrogen. Then, the seedlings were ground in a mortar with a pestle and the obtained powder was suspended in a medium containing 50 mmol/L Tris-HCl (pH 7.8, ratio 1:4; w/v), 0.1 mmol/L EDTA, 0.2% (v/v) Triton X-100 and 10% (v/v) glycerol. Homogenates were filtered through two layers of Miracloth and centrifuged at 27,000 g for 20 min. These supernatants were collected and utilized for the enzymatic *in vitro* assays.

Green pepper (*C. annuum* L.) fruit extracts were obtained in similar way but the resulting powder was re-suspended in 0.1 M Tris-HCl buffer (pH 8.0, ratio 1:1; w/v) containing 1 mmol/L EDTA, 0.1% (v/v) Triton X-100 and 10% (v/v) glycerol. Homogenates were also filtered through two layers of Miracloth and centrifuged at 27,000 g for 20 min. The supernatants were used for protein

enrichment by ammonium-sulfate [(NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>] to a quantity of 50% saturation. The obtained pellet was suspended in the same previous buffer and then loaded on a PD-10 desalting column containing Sephadex<sup>TM</sup> G-25 to eliminate the (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>.

### Enzyme activity assays in the presence of H<sub>2</sub>S

Catalase (EC 1.11.1.6) enzymatic activity was spectrophotometrically assayed by following the disappearance of H<sub>2</sub>O<sub>2</sub> at 240 nm (Aebi 1984). Glycolate oxidase (EC 1.1.3.1) activity was also determined spectrophotometrically by the formation of a glyoxylate-phenylhydrazine complex at 324 nm (Kerr and Groves 1975). NADH-dependent hydroxypyruvate reductase (HPR; EC 1.1.1.29) activity was assayed by monitoring the NADH oxidation at 340 nm (Schwitzguébel and Siegenthaler 1984).

For *in vitro* assays of persulfidation, the samples (*Arabidopsis* extracts, enriched pepper fruits and pure catalase from bovine liver) were previously treated in the absence and presence of sodium hydrosulfide (0.1, 0.5, 1.0, 2.0, 4.0, 5.0 and 10.0 mmol/L NaHS) as H<sub>2</sub>S donor for 60 min at 4°C in darkness (Aroca et al. 2015).

### Detection of hydrogen sulfide (H<sub>2</sub>S) in transgenic *Arabidopsis* seedlings expressing CFP-PTS1 and using CLSM technology

H<sub>2</sub>S was identified using 5 µM WSP-5 (Washington State Probe-5, Cayman Chemical) fluorescence probe dissolved in 10 mmol/L Tris-HCl (pH 7.4) buffer (Peng et al. 2014; Yu et al. 2014). The *Arabidopsis* seedlings were kept with this fluorescence probe (5 µM WSP-5 final concentration) in darkness conditions at 25°C for 1 h. Then, seedlings were washed twice in the same solution (10 mmol/L Tris-HCl, pH 7.4) for 15 min and placed on a microscopic slide. For examination, the confocal laser scanning microscope (Leica TCS SP5 II) was set up with the follow conditions: WSP-5 was excited at 502 nm and emission was collected at 525 nm and a 40 nm band pass width (490–530 nm); cyan fluorescent protein (CFP) was excited at 458 nm and emission was collected at 475 nm and a 40-nm band pass width (465–505 nm). As internal control, it was evaluated the potential overlap between the excitation and emission wavelengths of cyan fluorescent protein (CFP) with the fluorescent probe WSP-5 used to detect H<sub>2</sub>S. Additional controls were done to evaluate the efficiency and specificity of WSP5 to detect H<sub>2</sub>S in *Arabidopsis* samples. In this sense, 10-d-old *Arabidopsis* wild-type seedlings were

incubated with increasing concentrations of NaHS (0.1, 0.5 and 2 mmol/L) for 60 min at 25°C in the presence of 5 µM WSP5 and then observed by CLSM. In some cases, *Arabidopsis* seedlings were also incubated with 0.1 mmol/L hypotaurine, a specific H<sub>2</sub>S scavenger (Li et al. 2014; Shi et al. 2015) for 1 h at 25°C in the presence of 5 µM WSP5 and then observed by CLSM.

### Other assays

The measure of protein concentration was determined at 595 nm using the Bio-Rad protein assay (Hercules, CA) and a bovine serum albumin solution was used to prepare the standard curve. Relative fluorescence was quantified by using ImageJ software.

The results were the mean values ± standard errors (SE) obtained from a minimum of three independent biological replicates. The Student's t-test was used to determine the statistical significance between means. The half maximal inhibitory concentration (IC<sub>50</sub>) value represents the concentration at which a substance exerts half of its maximal inhibitory effect, and IC<sub>50</sub> for H<sub>2</sub>S was determined by on-line and easy-to-use software (<https://www.aatbio.com/tools/ic50-calculator/>).

## ACKNOWLEDGEMENTS

Mr. Carmelo Ruiz-Torres is deeply acknowledged to for his technical help. The “Centro de Instrumentación Científico-Técnica” (CICT) of University of Jaen is also recognized for the technical and human support (UJA, MINECO, Junta de Andalucía, FEDER). The project AGL2015-65104-P (ERDF grants co-financed by the Spanish Ministry of Economy and Competitiveness) supports the research in the laboratory of FJC, SGG, MAMV and JMP, and the grant BIO2015-66390-P for JBBs laboratory.

## AUTHOR CONTRIBUTIONS

F.J.C. designed the experiments, supervised the study, and wrote the manuscript. F.J.C. and J.B.B. carried out confocal scanning laser microscopy analysis. S.G-G and M.A.M-V performed enzymatic assays. F.J.C and J.M.P discussed the data and revised the manuscript. All authors read and approved the contents of this manuscript.

## REFERENCES

- Aebi H (1984) Catalase *in vitro*. **Method Enzymol** 105: 121–126
- Ali B, Gill RA, Yang S, Gill MB, Ali S, Rafiq MT, Zhou W. (2014) Hydrogen sulfide alleviates cadmium-induced morphophysiological and ultrastructural changes in *Brassica napus*. **Ecotoxicol Environ Saf** 110: 197–207
- Aroca Á, Serna A, Gotor C, Romero LC (2015) S-sulfhydration: A cysteine posttranslational modification in plant systems. **Plant Physiol** 168: 334–342
- Aroca A, Benito JM, Gotor C, Romero LC (2017) Persulfidation proteome reveals the regulation of protein function by hydrogen sulfide in diverse biological processes in *Arabidopsis*. **J Exp Bot** 68: 4915–4927
- Alvarez C, Calo L, Romero LC, García I, Gotor C (2010) An O-acetylserine(thiol)lyase homolog with L-cysteine desulfhydrase activity regulates cysteine homeostasis in *Arabidopsis*. **Plant Physiol** 152: 656–669
- Begara-Morales JC, López-Jaramillo FJ, Sánchez-Calvo B, Carreras A, Ortega-Muñoz M, Santoyo-González F, Corpas FJ, Barroso JB (2013) Vinyl sulfone silica: Application of an open preactivated support to the study of transnitrosylation of plant proteins by S-nitrosoglutathione. **BMC Plant Biol** 13: 61
- Bharwana SA, Ali S, Farooq MA, Ali B, Iqbal N, Abbas F, Ahmad MS (2014) Hydrogen sulfide ameliorates lead-induced morphological, photosynthetic, oxidative damages and biochemical changes in cotton. **Environ Sci Pollut Res Int**. 21: 717–731
- Birke H, De Kok LJ, Wirtz M, Hell R (2015) The role of compartment-specific cysteine synthesis for sulfur homeostasis during H<sub>2</sub>S exposure in *Arabidopsis*. **Plant Cell Physiol** 56: 358–367
- Calderwood A, Kopriva S (2014) Hydrogen sulfide in plants: From dissipation of excess sulfur to signaling molecule. **Nitric Oxide** 41: 72–78
- Chen J, Liu TW, Hu WJ, Simon M, Wang WH, Chen J, Liu X, Zheng HL (2014) Comparative proteomic analysis of differentially expressed proteins induced by hydrogen sulfide in *Spinacia oleracea* leaves. **PLoS ONE** 9: e105400
- Chen J, Wang WH, Wu FH, He EM, Liu X, Shangguan ZP, Zheng HL. (2015) Hydrogen sulfide enhances salt tolerance through nitric oxide-mediated maintenance of ion homeostasis in barley seedling roots. **Sci Rep** 5: 12516
- Chen Z, Chen M, Jiang M. (2017) Hydrogen sulfide alleviates mercury toxicity by sequestering it in roots or regulating reactive oxygen species productions in rice seedlings. **Plant Physiol Biochem** 111: 179–192
- Chaki M, Álvarez de Morales P, Ruiz C, Begara-Morales JC, Barroso JB, Corpas FJ, Palma JM (2015) Ripening of pepper (*Capsicum annuum*) fruit is characterized by an enhancement of protein tyrosine nitration. **Ann Bot** 116: 637–647
- Clark D, Durner J, Navarre DA, Klessig DF (2000) Nitric oxide inhibition of tobacco catalase and ascorbate peroxidase. **Mol Plant Microbe Interact** 13: 1380–1384
- Corpas FJ (2015) What is the role of hydrogen peroxide in plant peroxisomes? **Plant Biol (Stuttg)** 17: 1099–1103

- Corpas FJ, Barroso JB (2014) Peroxynitrite (ONOO<sup>-</sup>) is endogenously produced in *Arabidopsis* peroxisomes and is overproduced under cadmium stress. **Ann Bot** 113: 87–96
- Corpas FJ, Barroso JB (2015) Reactive sulfur species (RSS): Possible new players in the oxidative metabolism of plant peroxisomes. **Front Plant Sci** 6: 116
- Corpas FJ, Barroso JB (2017) Lead-induced stress, which triggers the production of nitric oxide (NO) and superoxide anion (O<sub>2</sub><sup>-</sup>) in *Arabidopsis* peroxisomes, affects catalase activity. **Nitric Oxide** 68: 103–110
- Corpas FJ, Barroso JB (2018) Peroxisomal plant metabolism: An update on nitric oxide, Ca<sup>2+</sup> and the NADPH recycling network. **J Cell Sci** 131: pii: jcs202978
- Corpas FJ, Barroso JB, Palma JM, Rodríguez-Ruiz M (2017) Plant peroxisomes: A nitro-oxidative cocktail. **Redox Biol** 11: 535–542
- Corpas FJ, González-Gordo G, Cañas A, Palma JM (2019) Nitric oxide (NO) and hydrogen sulfide (H<sub>2</sub>S) in plants: Which comes first? **J Exp Bot**. doi:10.1093/jxb/erz031
- Corpas FJ, Hayashi M, Mano S, Nishimura M, Barroso JB (2009) Peroxisomes are required for *in vivo* nitric oxide accumulation in the cytosol following salinity stress of *Arabidopsis* plants. **Plant Physiol** 151: 2083–2094
- Corpas FJ, Palma JM (2018) NO on/off in fruit ripening. **Plant Biol (Stuttg)** 20: 805–807
- Cuevasanta E, Möller MN, Alvarez B (2017) Biological chemistry of hydrogen sulfide and persulfides. **Arch Biochem Biophys** 617: 9–25
- de Freitas-Silva L, Rodríguez-Ruiz M, Houmani H, da Silva LC, Palma JM, Corpas FJ (2017) Glyphosate-induced oxidative stress in *Arabidopsis thaliana* affecting peroxisomal metabolism and triggers activity in the oxidative phase of the pentose phosphate pathway (OxPPP) involved in NADPH generation. **J Plant Physiol** 218: 196–205
- Filipovic MR (2015) Persulfidation (S-sulfhydration) and H<sub>2</sub>S. **Handb Exp Pharmacol** 230: 29–59
- Filipovic MR, Jovanovic VM (2017) More than just an intermediate: Hydrogen sulfide signalling in plants. **J Exp Bot** 68: 4733–4736
- Filipovic MR, Zivanovic J, Alvarez B, Banerjee R (2018) Chemical biology of H<sub>2</sub>S signaling through persulfidation. **Chem Rev** 118: 1253–1337
- Gotor C, Laureano-Marín AM, Moreno I, Aroca Á, García I, Romero LC (2015) Signaling in the plant cytosol: Cysteine or sulfide? **Amino Acids** 47: 2155–2164
- da Silva CJ, Batista Fontes EP, Modolo LV (2017) Salinity-induced accumulation of endogenous H<sub>2</sub>S and NO is associated with modulation of the antioxidant and redox defense systems in *Nicotiana tabacum* L. cv. Havana. **Plant Sci** 256: 148–159
- del Río LA, Corpas FJ, Sandalio LM, Palma JM, Gómez M, Barroso JB (2002) Reactive oxygen species, antioxidant systems and nitric oxide in peroxisomes. **J Exp Bot** 53: 1255–1272
- del Río LA (2015) ROS and RNS in plant physiology: An overview. **J Exp Bot** 66: 2827–2837
- Hancock JT, Whiteman M (2016) Hydrogen sulfide signaling: Interactions with nitric oxide and reactive oxygen species. **Ann N Y Acad Sci** 1365: 5–14
- Hänsch R, Lang C, Rennenberg H, Mendel RR (2007) Significance of plant sulfite oxidase. **Plant Biol (Stuttg)** 9: 589–595
- Hänsch R, Lang C, Riebeseel E, Lindigkeit R, Gessler A, Rennenberg H, Mendel RR (2006) Plant sulfite oxidase as novel producer of H<sub>2</sub>O<sub>2</sub>: Combination of enzyme catalysis with a subsequent non-enzymatic reaction step. **J Biol Chem** 281: 6884–6888
- Hänsch R, Mendel RR (2005) Sulfite oxidation in plant peroxisomes. **Photosynth Res** 86: 337–343
- Hu J, Baker A, Bartel B, Linka N, Mullen RT, Reumann S, Zolman BK (2012) Plant peroxisomes: Biogenesis and function. **Plant Cell** 24: 2279–2303
- Hu J, Huang X, Chen L, Sun X, Lu C, Zhang L, Wang Y, Zuo J (2015) Site-specific nitrosoproteomic identification of endogenously S-nitrosylated proteins in *Arabidopsis*. **Plant Physiol** 167: 1731–1746
- Hughes MN, Centelles MN, Moore KP (2009) Making and working with hydrogen sulfide: The chemistry and generation of hydrogen sulfide *in vitro* and its measurement *in vivo*: A review. **Free Radic Biol Med** 47: 1346–1353
- Iciek M, Kowalczyk-Pachel D, Bilska-Wilkosz A, Kwiecień I, Górny M, Włodek L (2015) S-sulfhydration as a cellular redox regulation. **Biosci Rep** 36: pii: e00304
- Jiménez A, Hernández JA, del Río LA, Sevilla F. (1997) Evidence for the presence of the ascorbate-glutathione cycle in mitochondria and peroxisomes of pea leaves. **Plant Physiol** 114: 275–284
- Ju Y, Fu M, Stokes E, Wu L, Yang G (2017) H<sub>2</sub>S-mediated protein S-sulfhydration: A prediction for its formation and regulation. **Molecules** 22: pii: E1334
- Khan MS, Haas FH, Samami AA, Gholami AM, Bauer A, Fellenberg K, Reichelt M, Hänsch R, Mendel RR, Meyer AJ, Wirtz M, Hell R (2010) Sulfite reductase defines a newly discovered bottleneck for assimilatory sulfate reduction and is essential for growth and development in *Arabidopsis thaliana*. **Plant Cell** 22: 1216–1231
- Kharbech O, Houmani H, Chaoui A, Corpas FJ (2017) Alleviation of Cr(VI)-induced oxidative stress in maize (*Zea mays* L.) seedlings by NO and H<sub>2</sub>S donors through differential organ-dependent regulation of ROS and NADPH-recycling metabolisms. **J Plant Physiol** 19: 71–80
- Kao YT, Gonzalez KL, Bartel B (2018) Peroxisome function, biogenesis, and dynamics in plants. **Plant Physiol** 176: 162–177
- Kerr MW, Groves D (1975) Purification and properties of glycollate oxidase from *Pisum sativum* leaves. **Phytochemistry** 14: 359–362
- Kimura H (2015) Signaling molecules: Hydrogen sulfide and polysulfide. **Antioxid Redox Signal** 22: 362–376
- Li YJ, Chen J, Xian M, Zhou LG, Han FX, Gan LJ, Shi ZQ (2014) *In site* bioimaging of hydrogen sulfide uncovers its pivotal

- role in regulating nitric oxide-induced lateral root formation. **PLoS ONE** 9: e90340
- Li ZG, Min X, Zhou ZH (2016) Hydrogen sulfide: A signal molecule in plant cross-adaptation. **Front Plant Sci** 7: 1621
- Li ZG, Yang SZ, Long WB, Yang GX, Shen ZZ (2013) Hydrogen sulphide may be a novel downstream signal molecule in nitric oxide-induced heat tolerance of maize (*Zea mays* L.) seedlings. **Plant Cell Environ** 36: 1564–1572
- Lisjak M, Teklic T, Wilson ID, Whiteman M, Hancock JT (2013) Hydrogen sulfide: Environmental factor or signalling molecule? **Plant Cell Environ** 36: 1607–1616
- Mathai JC, Missner A, Kügler P, Saparov SM, Zeidel ML, Lee JK, Pohl P (2009) No facilitator required for membrane transport of hydrogen sulfide. **Proc Natl Acad Sci USA** 106: 16633–16638
- Mhamdi A, Noctor G, Baker A (2012) Plant catalases: Peroxisomal redox guardians. **Arch Biochem Biophys** 525: 181–194
- Muñoz-Vargas MA, González-Gordo S, Cañas A, López-Jaramillo J, Palma JM, Corpas FJ (2018) Endogenous hydrogen sulfide (H<sub>2</sub>S) is up-regulated during sweet pepper (*Capsicum annuum* L.) fruit ripening. *In vitro* analyses show the inhibition of NADP-dependent isocitrate dehydrogenase activity by H<sub>2</sub>S and nitric oxide. **Nitric Oxide** 81: 36–45
- Mustafa AK, Gadalla MM, Sen N, Kim S, Mu W, Gazi SK, Barrow RK, Yang G, Wang R, Snyder SH (2009) H<sub>2</sub>S signals through protein S-sulfhydration. **Sci Signal** 2: ra72
- Neill SJ, Desikan R, Clarke A, Hurst RD, Hancock JT (2002) Hydrogen peroxide and nitric oxide as signalling molecules in plants. **J Exp Bot** 53: 1237–1247
- Nelson BK, Cai X, Nebenführ A (2007) A multicolored set of *in vivo* organelle markers for co-localization studies in *Arabidopsis* and other plants. **Plant J** 51: 1126–1136
- Nowak K, Luniak N, Witt C, Wüstefeld Y, Wachter A, Mendel RR, Hänsch R (2004) Peroxisomal localization of sulfite oxidase separates it from chloroplast-based sulfur assimilation. **Plant Cell Physiol** 45: 1889–1894
- Palma JM, Rodríguez-Ruiz M, Salvador González-Gordo S, Corpas FJ (2018) Interaction between antioxidants and reactive nitrogen species during pepper fruit ripening. **Free Rad Biol Med** 120: S6–S23
- Papenbrock J, Riemenschneider A, Kamp A, Schulz-Vogt HN, Schmidt A (2007) Characterization of cysteine-degrading and H<sub>2</sub>S-releasing enzymes of higher plants: From the field to the test tube and back. **Plant Biol** 9: 582–588
- Paul BD, Snyder SH (2015) H<sub>2</sub>S: A novel gasotransmitter that signals by sulfhydration. **Trends Biochem Sci** 40: 687–700
- Peek J, Christendat D (2015) The shikimate dehydrogenase family: Functional diversity within a conserved structural and mechanistic framework. **Arch Biochem Biophys** 566: 85–99
- Peng B, Chen W, Liu C, Rosser EW, Pacheco A, Zhao Y, Aguilar HC, Xian M (2014) Fluorescent probes based on nucleophilic substitution-cyclization for hydrogen sulfide detection and bioimaging. **Chemistry** 20: 1010–1016
- Riahi S, Rowley CN (2014) Why can hydrogen sulfide permeate cell membranes? **J Am Chem Soc** 136: 15111–15113
- Schwitzguébel JP, Siegenthaler PA (1984) Purification of peroxisomes and mitochondria from spinach leaf by percoll gradient centrifugation. **Plant Physiol** 75: 670–674
- Shi H, Ye T, Han N, Bian H, Liu X, Chan Z (2015) Hydrogen sulfide regulates abiotic stress tolerance and biotic stress resistance in *Arabidopsis*. **J Integr Plant Biol** 57: 628–640
- Su T, Wang P, Li H, Zhao Y, Lu Y, Dai P, Ren T, Wang X, Li X, Shao Q, Zhao D, Zhao Y, Ma C (2018) The *Arabidopsis* catalase triple mutant reveals important roles of catalases and peroxisome derived signaling in plant development. **J Integr Plant Biol** 60: 591–607
- Titov VY, Osipov AN (2017) Nitrite and nitroso compounds can serve as specific catalase inhibitors. **Redox Rep** 22: 91–97
- Tzin V, Galili G (2010) New insights into the shikimate and aromatic amino acids biosynthesis pathways in plants. **Mol Plant** 3: 956–972
- Veljovi-Jovanovic S, Oniki T, Takahama U (1998) Detection of monodehydro ascorbic acid radical in sulfite-treated leaves and mechanism of its formation. **Plant Cell Physiol** 39: 1203–1208
- Wang MJ, Cai WJ, Li N, Ding YJ, Chen Y, Zhu YC (2010) The hydrogen sulfide donor NaHS promotes angiogenesis in a rat model of hind limb ischemia. **Antioxid Redox Signal** 12: 1065–1077
- Yang M, Qin BP, Ma XL, Wang P, Li ML, Chen LL, Sun AQ, Wang ZL, Yin YP (2016) Foliar application of sodium hydrosulfide (NaHS), a hydrogen sulfide (H<sub>2</sub>S) donor, can protect seedlings against heat stress in wheat (*Triticum aestivum* L.). **J Integr Agric** 15: 2745–2758
- Yamasaki H, Cohen MF (2016) Biological consilience of hydrogen sulfide and nitric oxide in plants: Gases of primordial earth linking plant, microbial and animal physiologies. **Nitric Oxide** 55–56: 91–100
- Yu F, Han X, Chen L (2014) Fluorescent probes for hydrogen sulfide detection and bioimaging. **Chem Commun (Camb)** 50: 12234–12249
- Zhang P, Luo Q, Wang R, Xu J (2017) Hydrogen sulfide toxicity inhibits primary root growth through the ROS-NO pathway. **Sci Rep** 7: 868
- Zhang H, Tan ZQ, Hu LY, Wang SH, Luo JP, Jones RL (2010) Hydrogen sulfide alleviates aluminum toxicity in germinating wheat seedlings. **J Integr Plant Biol** 52: 556–567
- Zhang D, Macinkovic I, Devarie-Baez NO, Pan J, Park CM, Carroll KS, Filipovic MR, Xian M (2014) Detection of protein S-sulfhydration by a tag-switch technique. **Angew Chem** 53: 575–581

## SUPPORTING INFORMATION

Additional Supporting Information may be found online in the supporting information tab for this article: <http://onlinelibrary.wiley.com/doi/10.1111/jipb.12779/suppinfo>  
**Figure S1. Control to evaluate the potential overlap between the excitation and emission wavelengths of**

**cyan fluorescent protein (CFP) with the fluorescent probe WSP-5**

Representative images illustrating the CLSM *in vivo* detection of peroxisomes (green) in root tips of transgenic *Arabidopsis* 10-d-old seedlings expressing CFP-PTS1 without the presence of WSP-5, fluorescent probe used to detect H<sub>2</sub>S, and observed under two conditions. **(A)** Shows fluorescence punctuates

(green) attributable to CFP-PTS1 (excitation 458 nm; emission 475 nm) indicating the localization of peroxisomes. **(B)** Shows the absence of any fluorescence in the same area observed in **(A)**, respectively, without any fluorescence probe WSP-5 (negative control) using the wavelength conditions to detect H<sub>2</sub>S with WSP-5 (excitation 502 nm and emission 525 nm).



Scan using WeChat with your smartphone to view JIPB online



Scan with iPhone or iPad to view JIPB online